

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Total Unintentional Injuries/Violence	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN8: Percentage of students who rarely or never wore a bicycle helmet (among students who had ridden a bicycle during the 12 months before the survey)	36.8 (34.2-39.4)	60.1 (57.9-62.3)	77.7 (74.6-80.5)	80.8 (74.6-85.8)	Yes
QN9: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)	3.6 (2.7-5.0)	6.5 (5.7-7.3)	11.9 (10.5-13.4)	29.4 (24.8-34.4)	Yes
QN10: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle one or more times during the 30 days before the survey)	12.9 (11.2-14.7)	15.5 (13.9-17.2)	16.3 (14.4-18.5)	33.5 (29.3-38.1)	Yes
QN11: Percentage of students who drove when drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	3.9 (3.0-5.0)	5.6 (4.7-6.5)	6.3 (5.0-7.8)	24.5 (19.1-30.9)	Yes
QN12: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	42.2 (39.3-45.1)	45.7 (42.8-48.5)	41.1 (36.8-45.6)	45.5 (38.4-52.8)	Yes
QN16: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	2.9 (2.2-3.8)	4.5 (3.8-5.2)	7.6 (6.3-9.2)	13.6 (10.5-17.4)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Total Unintentional Injuries/Violence	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN19: Percentage of students who were injured in a physical fight (one or more times during the 12 months before the survey; injuries had to be treated by a doctor or nurse)	2.1 (1.6-2.9)	3.0 (2.5-3.5)	5.7 (4.6-7.1)	12.9 (10.1-16.2)	Yes
QN20: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	2.9 (2.3-3.7)	4.7 (4.1-5.3)	9.2 (8.0-10.5)	22.2 (18.1-26.9)	Yes
QN21: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	3.8 (2.8-5.1)	6.0 (5.2-6.8)	7.9 (6.6-9.5)	15.4 (11.7-20.0)	Yes
QN22: Percentage of students who experienced physical dating violence (one or more times during the 12 months before the survey, including being hit, slammed into something, or injured with an object or weapon on purpose by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	5.1 (4.0-6.6)	6.9 (5.9-8.0)	10.3 (8.4-12.5)	16.1 (12.4-20.7)	Yes
QN23: Percentage of students who experienced sexual dating violence (one or more times during the 12 months before the survey, including kissing, touching, or being physically forced to have sexual intercourse when they did not want to by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	9.1 (7.2-11.4)	11.9 (10.3-13.6)	11.9 (10.3-13.8)	20.1 (15.5-25.6)	Yes
QN24: Percentage of students who were bullied on school property (during the 12 months before the survey)	15.9 (14.2-17.8)	22.0 (20.5-23.5)	28.1 (25.1-31.3)	35.7 (31.2-40.5)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Total Unintentional Injuries/Violence	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN25: Percentage of students who were electronically bullied (including being bullied through e-mail, chat rooms, instant messaging, websites, or texting during the 12 months before the survey)	12.9 (11.4-14.6)	19.5 (18.0-21.1)	21.5 (18.7-24.6)	29.6 (25.2-34.4)	Yes
QN26: Percentage of students who felt sad or hopeless (almost every day for 2 or more weeks in a row so that they stopped doing some usual activities during the 12 months before the survey)	19.9 (18.0-22.0)	27.5 (25.6-29.4)	34.5 (32.0-37.2)	44.7 (39.7-49.8)	Yes
QN27: Percentage of students who seriously considered attempting suicide (during the 12 months before the survey)	10.1 (8.6-11.8)	14.9 (13.6-16.3)	20.3 (18.1-22.6)	34.3 (29.7-39.1)	Yes
QN29: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	3.0 (2.5-3.6)	5.9 (5.1-6.9)	10.8 (9.1-12.7)	22.3 (17.6-27.9)	Yes
QN30: Percentage of students who attempted suicide that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	0.8 (0.6-1.1)	1.9 (1.4-2.4)	4.7 (3.4-6.3)	9.7 (6.7-14.0)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Total  
Tobacco Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN32: Percentage of students who smoked a whole cigarette before age 13 years (for the first time)	2.2 (1.7-3.0)	3.9 (3.4-4.4)	8.8 (7.6-10.2)	22.6 (18.3-27.6)	Yes
QN33: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	3.2 (2.7-3.8)	7.7 (6.9-8.7)	15.8 (14.1-17.7)	32.5 (27.5-37.9)	Yes
QNFRCIG: Percentage of students who currently frequently smoked cigarettes (on 20 or more days during the 30 days before the survey)	0.9 (0.7-1.3)	2.7 (2.3-3.2)	6.4 (5.2-7.9)	16.4 (12.8-20.9)	Yes
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	0.7 (0.5-1.0)	1.9 (1.5-2.3)	4.1 (3.3-5.1)	13.8 (10.3-18.1)	Yes
QN34: Percentage of students who smoked more than 10 cigarettes per day (during the 30 days before the survey among students who currently smoked cigarettes on the days they smoked)	12.5 (7.9-19.2)	7.6 (5.3-10.7)	8.5 (5.8-12.2)	26.6 (19.3-35.3)	Yes
QN35: Percentage of students who usually obtained their own cigarettes by buying them in a store or gas station (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	12.4 (7.2-20.6)	9.4 (6.6-13.3)	14.3 (10.5-19.3)	22.4 (15.8-30.7)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Total  
Tobacco Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNCIGINT: Percentage of students who usually obtained their own cigarettes by buying on the internet (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	1.9 (0.6-5.7)	1.4 (0.5-3.8)	1.5 (0.5-4.7)	3.1 (1.1-8.2)	No
QN36: Percentage of students who tried to quit smoking cigarettes (among students who currently smoked cigarettes during the 12 months before the survey)	41.1 (33.3-49.3)	41.6 (36.6-46.8)	55.5 (49.7-61.1)	44.5 (36.3-53.0)	Yes
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, or dip on at least 1 day during the 30 days before the survey)	2.4 (1.8-3.2)	5.0 (4.3-5.8)	9.4 (7.8-11.3)	21.5 (16.7-27.1)	Yes
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars on at least 1 day during the 30 days before the survey)	5.9 (4.9-7.0)	11.0 (9.7-12.5)	15.4 (13.3-17.8)	26.5 (22.1-31.4)	Yes
QN40: Percentage of students who currently used electronic vapor products (e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens such as blu, NJOY, or Starbuzz on at least 1 day during the 30 days before the survey)	13.4 (11.8-15.1)	26.4 (24.6-28.3)	36.1 (32.9-39.4)	52.3 (46.7-57.8)	Yes
QNTOB4: Percentage of students who currently used tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	16.9 (14.9-19.1)	31.5 (29.5-33.7)	43.8 (40.2-47.4)	62.4 (57.1-67.5)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Total  
Tobacco Use

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNTOB3: Percentage of students who currently used cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	8.2 (7.0-9.5)	16.6 (15.0-18.3)	27.7 (24.8-30.8)	44.0 (38.1-49.9)	Yes
QNTOB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	7.5 (6.4-8.8)	15.3 (13.7-16.9)	24.1 (21.6-26.7)	40.1 (34.8-45.7)	Yes
QNNOTOB4: Percentage of students who did not currently use tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	83.8 (81.7-85.7)	69.3 (67.2-71.4)	57.3 (53.6-60.9)	39.2 (33.9-44.6)	Yes
QNNOTOB3: Percentage of students who did not currently use cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	92.5 (91.3-93.5)	84.1 (82.4-85.6)	73.0 (69.9-75.9)	57.7 (51.6-63.5)	Yes
QNNOTOB2: Percentage of students who did not currently smoke cigarettes or cigars (on at least 1 day during the 30 days before the survey)	93.1 (92.0-94.1)	85.4 (83.8-86.9)	76.5 (73.9-79.0)	61.2 (55.6-66.6)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Total Alcohol/Other Drug Use	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN42: Percentage of students who drank alcohol before age 13 years (for the first time other than a few sips)	5.7 (4.6-6.9)	9.6 (8.5-10.7)	14.9 (13.2-16.9)	30.1 (25.1-35.5)	Yes
QN43: Percentage of students who currently drank alcohol (at least one drink of alcohol on at least 1 day during the 30 days before the survey)	21.6 (19.5-23.7)	32.9 (30.3-35.7)	35.6 (32.8-38.5)	47.2 (40.7-53.7)	Yes
QN44: Percentage of students who drank five or more drinks of alcohol in a row (within a couple of hours on at least 1 day during the 30 days before the survey)	11.5 (9.9-13.2)	17.4 (15.7-19.3)	21.4 (19.1-23.8)	31.7 (26.5-37.3)	Yes
QN46: Percentage of students who usually obtained the alcohol they drank by someone giving it to them (among students who currently drank alcohol)	51.0 (47.3-54.6)	41.9 (39.2-44.6)	37.9 (33.2-42.8)	31.3 (23.3-40.7)	Yes
QN48: Percentage of students who tried marijuana before age 13 years (for the first time)	2.0 (1.6-2.5)	5.1 (4.1-6.4)	10.6 (9.0-12.4)	21.6 (17.4-26.6)	Yes
QN49: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	11.4 (10.1-12.8)	22.7 (20.6-25.0)	34.2 (31.1-37.5)	41.4 (36.1-46.8)	Yes
QN50: Percentage of students who ever used cocaine (any form of cocaine, such as powder, crack, or freebase, one or more times during their life)	2.4 (1.8-3.1)	3.6 (3.0-4.2)	7.8 (6.5-9.4)	17.5 (13.7-21.9)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Total Alcohol/Other Drug Use	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	3.5 (2.8-4.3)	5.3 (4.7-6.0)	8.5 (7.4-9.8)	22.3 (18.3-27.0)	Yes
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China white," one or more times during their life)	1.3 (0.9-2.0)	1.5 (1.1-1.8)	2.6 (2.0-3.4)	10.6 (8.0-13.8)	Yes
QN53: Percentage of students who ever used methamphetamines (also called "speed," "crystal," "crank," or "ice," one or more times during their life)	1.5 (1.1-2.2)	1.5 (1.2-1.9)	2.9 (2.3-3.7)	9.0 (6.6-12.0)	Yes
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	1.8 (1.4-2.3)	3.8 (3.2-4.6)	7.0 (5.7-8.6)	13.8 (10.6-17.8)	Yes
QN55: Percentage of students who ever used synthetic marijuana (also called "K2", "Spice", "fake weed", "King Kong", "Yucatan Fire", "Skunk", or "Moon Rocks", one or more times during their life)	3.5 (2.9-4.4)	8.2 (7.2-9.4)	14.8 (12.7-17.1)	27.7 (23.0-32.9)	Yes
QN57: Percentage of students who ever took prescription drugs without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during their life)	7.2 (6.3-8.3)	13.1 (11.8-14.5)	18.3 (16.3-20.5)	36.3 (31.5-41.4)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.



## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Total Sexual Behaviors	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
<b>Health Risk Behavior</b>					
QN60: Percentage of students who ever had sexual intercourse	27.4 (24.8-30.2)	40.9 (38.2-43.6)	49.2 (45.9-52.5)	62.3 (56.6-67.6)	Yes
QN61: Percentage of students who had sexual intercourse before age 13 years (for the first time)	1.5 (1.1-2.1)	1.7 (1.4-2.1)	3.5 (2.7-4.5)	10.9 (7.9-15.0)	Yes
QN63: Percentage of students who were currently sexually active (sexual intercourse with at least one person during the 3 months before the survey)	21.9 (19.4-24.6)	32.7 (30.4-35.0)	38.9 (35.8-42.1)	47.8 (41.6-54.2)	Yes
QN64: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	13.9 (11.1-17.2)	14.9 (13.1-17.0)	23.9 (20.7-27.4)	36.0 (27.9-45.0)	Yes
QN65: Percentage of students who used a condom (during last sexual intercourse among students who were currently sexually active)	67.2 (62.8-71.2)	60.7 (58.0-63.4)	59.0 (54.1-63.8)	43.9 (33.4-54.9)	Yes
QN66: Percentage of students who used birth control pills (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	46.3 (41.3-51.4)	32.8 (29.9-35.8)	26.0 (22.3-30.0)	23.4 (15.5-33.7)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Total Sexual Behaviors	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
<b>Health Risk Behavior</b>					
QNIUDIMP: Percentage of students who used an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	3.7 (2.7-5.1)	5.6 (4.3-7.1)	5.4 (4.0-7.3)	3.6 (1.9-6.8)	Yes
QNSHPARG: Percentage of students who used a shot (e.g., Depo-Provera), patch (e.g., OrthoEvra), or birth control ring (e.g., NuvaRing) (during last sexual intercourse among students who were currently sexually active)	2.9 (2.0-4.2)	4.2 (3.3-5.4)	5.2 (3.1-8.5)	1.8 (0.7-4.3)	Yes
QNOTHHPL: Percentage of students who used birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	52.9 (47.9-57.9)	42.6 (39.6-45.7)	36.5 (31.8-41.5)	28.8 (20.7-38.5)	Yes
QNDUALBC: Percentage of students who used both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	30.0 (25.0-35.5)	16.1 (14.3-18.2)	12.5 (9.8-15.8)	7.2 (4.3-11.7)	Yes
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy (during last sexual intercourse among students who were currently sexually active)	3.9 (2.8-5.4)	6.4 (5.3-7.8)	7.3 (5.6-9.4)	19.4 (13.4-27.3)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Total Dietary Behaviors	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
<b>Health Risk Behavior</b>					
QNOBESE: Percentage of students who were obese ( $\geq$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart <sup>§</sup> )	6.7 (5.8-7.7)	12.4 (11.3-13.5)	17.2 (15.3-19.1)	23.1 (18.1-29.0)	Yes
QNOWT: Percentage of students who were overweight ( $\geq$ 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart <sup>§</sup> )	12.8 (11.2-14.5)	14.3 (12.8-15.9)	17.1 (14.3-20.4)	16.7 (13.3-20.6)	Yes
QN70: Percentage of students who were trying to lose weight	41.7 (38.7-44.6)	45.2 (43.3-47.1)	45.7 (42.8-48.6)	44.9 (38.0-51.9)	Yes
QN77: Percentage of students who did not drink a can, bottle, or glass of soda or pop (not including diet soda or diet pop, during the 7 days before the survey)	47.8 (45.1-50.5)	34.7 (32.9-36.6)	27.5 (24.9-30.3)	21.1 (17.0-25.9)	Yes
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	6.1 (5.2-7.3)	12.2 (10.8-13.6)	20.6 (18.5-22.9)	31.2 (26.5-36.4)	Yes
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	3.2 (2.6-4.0)	8.2 (7.2-9.3)	16.6 (14.7-18.8)	26.7 (22.2-31.8)	Yes
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	1.5 (1.1-1.8)	4.2 (3.5-5.0)	8.9 (7.7-10.4)	16.3 (12.9-20.4)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school,  $p < 0.05$ .

† Confidence interval.

§ Based on reference data from the 2000 CDC Growth Charts.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Total Physical Activity	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
<b>Health Risk Behavior</b>					
QN80: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	52.6 (49.9-55.3)	46.3 (43.6-49.1)	42.9 (39.7-46.1)	36.6 (30.7-42.8)	Yes
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	10.5 (9.1-12.1)	12.0 (10.8-13.3)	16.6 (14.7-18.7)	24.8 (20.1-30.1)	Yes
QNPA7DAY: Percentage of students who were physically active at least 60 minutes per day on all 7 days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time (during the 7 days before the survey)	21.2 (19.2-23.3)	22.9 (21.0-24.9)	22.1 (19.3-25.2)	19.4 (15.8-23.6)	Yes
QN81: Percentage of students who watched television 3 or more hours per day (on an average school day)	13.0 (11.5-14.6)	19.7 (18.3-21.1)	24.4 (21.8-27.1)	30.5 (26.1-35.2)	Yes
QN82: Percentage of students who played video or computer games or used a computer 3 or more hours per day (for something that was not school work on an average school day)	31.3 (29.1-33.7)	39.0 (37.2-40.9)	46.0 (43.3-48.7)	53.1 (46.6-59.6)	Yes
QN88: Percentage of students who had 8 or more hours of sleep (on an average school night)	29.2 (27.0-31.5)	29.4 (27.2-31.7)	24.9 (22.6-27.3)	21.3 (17.5-25.6)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Total Other	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school					Significant Association*
	A's	B's	C's	D's/F's		
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI		
	<b>Health Risk Behavior</b>					
	QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work during the 12 months before the survey)	90.1 (88.8-91.2)	83.9 (82.4-85.4)	76.9 (74.6-79.1)	63.2 (57.8-68.3)	Yes
	QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma	21.7 (19.7-23.8)	21.3 (20.1-22.6)	23.7 (21.3-26.2)	25.7 (20.6-31.7)	No

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Total  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN90: Percentage of students who never or rarely wear a seat belt when driving (among students who drive a car)	3.4 (2.6-4.4)	5.9 (5.1-6.8)	11.9 (10.0-14.1)	27.3 (20.9-34.8)	Yes
QN91: Percentage of students who have experienced an unwanted sexual advance because of another student's drinking (one or more times during the 12 months before the survey)	4.6 (3.7-5.6)	4.7 (4.0-5.4)	7.2 (5.6-9.2)	11.8 (8.2-16.8)	Yes
QN92: Percentage of students who ever saw or heard adults in their home slap, hit, kick, punch, or hurt each other	10.7 (9.0-12.7)	13.1 (12.0-14.4)	18.8 (16.2-21.8)	27.1 (22.5-32.2)	Yes
QN93: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)	13.2 (11.9-14.7)	18.3 (16.3-20.3)	23.5 (20.6-26.6)	33.4 (28.5-38.8)	Yes
QN94: Percentage of students who ever came to school high on marijuana (during their life)	6.7 (5.6-8.1)	14.8 (13.4-16.3)	25.7 (23.3-28.2)	36.9 (31.4-42.8)	Yes
QN95: Percentage of students who currently took a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during the 30 days before the survey)	3.1 (2.6-3.8)	6.2 (5.4-7.1)	8.7 (7.3-10.2)	23.5 (19.3-28.4)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Total  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN96: Percentage of students who ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight (during the 30 days before the survey)	39.2 (35.7-42.7)	37.6 (35.8-39.5)	39.3 (36.1-42.5)	38.9 (33.1-45.1)	Yes
QN97: Percentage of students who will play on their school's sports team one or more seasons (during the school year of the survey)	67.6 (65.2-69.8)	61.2 (58.7-63.6)	47.7 (43.7-51.6)	31.2 (26.0-37.0)	Yes
QN98: Percentage of students who used an indoor tanning device (such as a sunlamp, sunbed, or tanning booth [not including getting a spray-on tan], one or more times during the 12 months before the survey)	4.6 (3.6-5.7)	6.3 (5.4-7.4)	6.0 (4.8-7.5)	8.3 (6.2-11.2)	Yes
QN99: Percentage of students who saw a doctor or nurse (for a check-up or physical exam when they were not sick or injured during the 12 months before the survey)	85.9 (84.4-87.2)	80.0 (78.2-81.6)	74.4 (71.9-76.8)	67.6 (62.0-72.7)	Yes
QN100: Percentage of students who have someone in their family (a parent, brother, or sister) who is currently in the military (air Force, Army, Marines, National Guard, Navy, or Reserves)	10.6 (9.2-12.3)	13.6 (12.4-14.9)	22.4 (19.0-26.2)	22.1 (17.3-27.7)	Yes
QN101: Percentage of students who reported either of their parents or other adults in their family have been in jail or in prison (during the 12 months before the survey)	5.0 (4.0-6.3)	8.2 (7.3-9.1)	13.8 (12.0-15.7)	22.1 (18.3-26.5)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Total  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN102: Percentage of students who strongly agree or agree that their parents or other adults in their family have clear rules and consequences for their behavior	83.2 (81.4-84.8)	77.1 (75.2-78.9)	71.8 (69.4-74.0)	52.8 (47.7-57.8)	Yes
QN103: Percentage of students who went hungry most of the time or always because there was not enough food in their home	1.2 (0.8-1.9)	2.0 (1.5-2.7)	2.9 (2.1-4.1)	9.0 (6.7-12.1)	Yes
QN104: Percentage of students who have ever lived with someone who had a problem with alcohol or drugs	20.9 (18.9-23.1)	30.0 (28.1-32.0)	42.7 (39.3-46.2)	52.4 (46.9-57.8)	Yes
QN105: Percentage of students who have talked with at least one of their parents or guardians about the dangers of tobacco, alcohol, or drug use (during the 12 months before the survey)	55.2 (52.3-57.9)	49.8 (47.3-52.3)	46.4 (42.7-50.1)	41.8 (36.3-47.5)	Yes
QN106: Percentage of students who recall hearing, reading, or seeing a message about avoiding alcohol or other illegal drugs (during the 12 months before the survey)	85.7 (84.3-87.1)	80.4 (78.8-81.9)	74.0 (71.6-76.2)	69.4 (64.6-73.9)	Yes
QN107: Percentage of students who would most likely talk with their parent or other adult family member about their feelings (among students who report having felt sad, empty, hopeless, angry, or anxious)	26.6 (23.9-29.4)	18.9 (17.2-20.7)	15.1 (12.4-18.2)	9.6 (6.9-13.1)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Total  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN108: Percentage of students who performed organized community service as a non-paid volunteer (such as serving meals to the elderly, picking up litter, helping out at a hospital, or building homes for the poor, one or more times during the 30 days before the survey)	56.4 (54.0-58.7)	41.0 (38.9-43.1)	28.4 (25.5-31.5)	27.2 (21.1-34.3)	Yes
QN109: Percentage of students who participate in activities run by community groups (including activities run by 4-H, Boys and Girls Club, YMCA, sports clubs, or church groups, for one or more hours during an average week when they are in school)	49.7 (46.4-53.0)	35.7 (33.5-38.0)	23.7 (21.5-26.2)	17.2 (13.6-21.5)	Yes
QN110: Percentage of students who think people are at great risk harming themselves if they smoke one or more packs of cigarettes per day (physically or in other ways)	80.1 (78.2-81.9)	70.9 (69.1-72.5)	60.1 (56.7-63.5)	53.4 (47.7-59.1)	Yes
QN111: Percentage of students who think people are at great risk harming themselves if they have one or two drinks of alcohol nearly every day (physically or in other ways)	31.1 (28.8-33.4)	25.9 (24.3-27.6)	23.5 (20.2-27.1)	17.4 (13.6-21.9)	Yes
QN112: Percentage of students who think people are at great risk of harming themselves (physically or in other ways) if they have five or more drinks of alcohol once or twice a week (beer, wine, or liquor)	49.9 (47.2-52.6)	39.6 (37.9-41.4)	31.9 (29.4-34.6)	27.0 (22.7-31.6)	Yes
QN113: Percentage of students who think people are at great risk of harming themselves if they use marijuana once or twice a week (physically or in other ways)	24.5 (22.4-26.8)	17.3 (15.4-19.4)	11.0 (9.5-12.7)	9.1 (6.0-13.6)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Total Site-Added	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN114: Percentage of students who think people are at great risk of harming themselves if they take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	70.0 (67.4-72.5)	63.5 (61.3-65.6)	54.6 (51.5-57.6)	48.5 (42.5-54.6)	Yes
QN115: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to smoke tobacco	78.2 (75.7-80.6)	66.0 (64.2-67.8)	52.8 (49.8-55.8)	39.9 (32.9-47.3)	Yes
QN116: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to have one or two drinks of alcoholic beverage nearly every day (beer, wine or liquor)	69.7 (67.6-71.7)	56.0 (53.8-58.2)	46.1 (42.3-49.9)	35.2 (29.7-41.2)	Yes
QN117: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to smoke marijuana	55.8 (52.0-59.4)	41.5 (38.8-44.2)	30.2 (27.0-33.6)	21.5 (17.4-26.3)	Yes
QN118: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	88.5 (86.9-89.9)	82.1 (80.6-83.6)	76.2 (73.9-78.3)	61.1 (54.4-67.4)	Yes
QN119: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to smoke tobacco	95.5 (94.3-96.4)	90.9 (89.8-92.0)	85.7 (83.2-87.8)	74.5 (69.4-79.0)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Total Site-Added	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN120: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to have one or two drinks of an alcoholic beverage nearly every day (beer, wine or liquor)	93.5 (92.3-94.6)	90.0 (88.8-91.1)	87.0 (85.3-88.6)	74.1 (68.9-78.7)	Yes
QN121: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to smoke marijuana	91.5 (90.4-92.5)	82.5 (80.8-84.1)	75.6 (73.1-78.0)	64.8 (59.6-69.6)	Yes
QN122: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	97.2 (96.5-97.7)	95.5 (94.8-96.2)	94.0 (92.6-95.1)	85.6 (81.0-89.2)	Yes
QN123: Percentage of students who approve or strongly approve of someone their age having one or two drinks of alcohol nearly every day (beer, wine, or liquor)	5.1 (3.8-7.0)	6.8 (6.0-7.7)	12.5 (10.7-14.5)	19.9 (16.2-24.3)	Yes
QN124: Percentage of students who think it would be very easy for them to get some cigarettes if they wanted to	26.8 (24.2-29.5)	37.4 (35.1-39.9)	45.4 (41.7-49.1)	53.6 (48.4-58.6)	Yes
QN125: Percentage of students who think it would be very easy for them to get some alcohol if they wanted to (beer, wine, or liquor)	34.8 (32.4-37.3)	38.6 (36.6-40.7)	39.5 (36.5-42.5)	44.7 (39.6-50.0)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Total  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN126: Percentage of students who think it would be very easy for them to get some marijuana if they wanted to	31.1 (28.9-33.3)	43.7 (41.6-45.7)	50.5 (47.2-53.8)	61.0 (56.1-65.7)	Yes
QN127: Percentage of students who think it would be very easy for them to get some prescription drug without a doctor's prescription if they wanted to (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	10.2 (8.8-11.9)	13.9 (12.5-15.5)	20.0 (17.3-23.0)	27.2 (22.8-32.0)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.



## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Male Unintentional Injuries/Violence	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN19: Percentage of students who were injured in a physical fight (one or more times during the 12 months before the survey; injuries had to be treated by a doctor or nurse)	3.7 (2.5-5.5)	3.6 (3.0-4.3)	5.7 (4.5-7.1)	12.2 (9.0-16.2)	Yes
QN20: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	5.4 (3.9-7.3)	5.8 (4.9-7.0)	11.1 (9.5-12.9)	23.3 (18.3-29.2)	Yes
QN21: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	1.7 (1.2-2.5)	2.0 (1.5-2.6)	2.6 (1.8-3.7)	9.1 (5.4-14.8)	Yes
QN22: Percentage of students who experienced physical dating violence (one or more times during the 12 months before the survey, including being hit, slammed into something, or injured with an object or weapon on purpose by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	5.1 (3.4-7.7)	4.5 (3.3-5.9)	6.6 (4.6-9.4)	11.3 (7.9-15.9)	Yes
QN23: Percentage of students who experienced sexual dating violence (one or more times during the 12 months before the survey, including kissing, touching, or being physically forced to have sexual intercourse when they did not want to by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	4.2 (3.1-5.5)	4.9 (3.7-6.4)	5.7 (4.3-7.5)	12.2 (7.3-19.5)	Yes
QN24: Percentage of students who were bullied on school property (during the 12 months before the survey)	11.6 (9.7-13.9)	14.7 (13.1-16.6)	20.5 (17.9-23.4)	29.7 (24.4-35.7)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Male Unintentional Injuries/Violence		Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
Health Risk Behavior	A's	B's	C's	D's/F's		
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI		
QN25: Percentage of students who were electronically bullied (including being bullied through e-mail, chat rooms, instant messaging, websites, or texting during the 12 months before the survey)	7.1 (5.6-9.0)	10.6 (8.7-12.9)	13.4 (10.4-17.1)	22.8 (17.7-28.9)	Yes	
QN26: Percentage of students who felt sad or hopeless (almost every day for 2 or more weeks in a row so that they stopped doing some usual activities during the 12 months before the survey)	11.8 (9.8-14.2)	16.1 (14.1-18.4)	22.4 (19.8-25.1)	34.7 (28.8-41.1)	Yes	
QN27: Percentage of students who seriously considered attempting suicide (during the 12 months before the survey)	6.4 (5.0-8.1)	9.2 (7.8-10.8)	13.1 (11.2-15.3)	27.0 (21.6-33.3)	Yes	
QN29: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	1.6 (1.1-2.3)	3.3 (2.3-4.7)	6.2 (5.0-7.7)	17.1 (11.6-24.4)	Yes	
QN30: Percentage of students who attempted suicide that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	0.5 (0.2-1.0)	0.9 (0.6-1.4)	2.0 (1.3-3.0)	7.6 (4.3-13.0)	Yes	

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male  
Tobacco Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN32: Percentage of students who smoked a whole cigarette before age 13 years (for the first time)	3.6 (2.4-5.3)	4.2 (3.5-5.0)	9.8 (8.2-11.6)	24.0 (18.5-30.5)	Yes
QN33: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	4.1 (3.2-5.3)	7.6 (6.4-8.9)	14.5 (12.6-16.7)	31.2 (25.2-37.8)	Yes
QNFRCIG: Percentage of students who currently frequently smoked cigarettes (on 20 or more days during the 30 days before the survey)	1.3 (0.8-2.0)	2.7 (2.1-3.4)	6.5 (5.2-8.1)	15.8 (11.6-21.0)	Yes
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	1.0 (0.6-1.7)	1.9 (1.4-2.5)	4.3 (3.2-5.8)	13.4 (9.6-18.4)	Yes
QN34: Percentage of students who smoked more than 10 cigarettes per day (during the 30 days before the survey among students who currently smoked cigarettes on the days they smoked)	17.6 (10.5-28.2)	9.3 (5.8-14.5)	12.4 (8.3-18.2)	26.7 (19.0-36.2)	Yes
QN35: Percentage of students who usually obtained their own cigarettes by buying them in a store or gas station (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	16.1 (8.1-29.6)	12.5 (7.7-19.7)	20.0 (14.2-27.5)	28.6 (19.3-40.1)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male  
Tobacco Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNCIGINT: Percentage of students who usually obtained their own cigarettes by buying on the internet (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	2.5 (0.6-9.5)	2.3 (0.7-7.0)	1.7 (0.4-7.0)	1.3 (0.3-6.0)	No
QN36: Percentage of students who tried to quit smoking cigarettes (among students who currently smoked cigarettes during the 12 months before the survey)	40.8 (29.3-53.4)	43.5 (35.8-51.5)	49.0 (41.8-56.3)	40.0 (30.1-50.7)	No
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, or dip on at least 1 day during the 30 days before the survey)	5.0 (3.7-6.9)	8.5 (7.3-10.0)	13.1 (10.8-15.7)	25.6 (19.5-32.9)	Yes
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars on at least 1 day during the 30 days before the survey)	10.7 (8.4-13.4)	15.3 (13.4-17.5)	18.6 (16.0-21.5)	28.7 (23.2-35.0)	Yes
QN40: Percentage of students who currently used electronic vapor products (e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens such as blu, NJOY, or Starbuzz on at least 1 day during the 30 days before the survey)	13.3 (11.4-15.5)	25.6 (23.1-28.2)	36.1 (32.4-40.0)	50.8 (43.9-57.8)	Yes
QNTOB4: Percentage of students who currently used tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	19.8 (17.0-22.9)	32.7 (29.9-35.6)	45.1 (41.0-49.1)	60.7 (54.0-67.0)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male  
Tobacco Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNTOB3: Percentage of students who currently used cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	13.5 (11.0-16.3)	20.6 (18.5-23.0)	30.5 (26.8-34.4)	45.1 (37.8-52.7)	Yes
QNTOB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	12.1 (9.8-14.9)	18.5 (16.3-20.8)	25.4 (22.4-28.6)	40.1 (33.5-47.1)	Yes
QNNOTOB4: Percentage of students who did not currently use tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	81.6 (78.7-84.2)	68.5 (65.6-71.2)	55.9 (51.8-60.0)	40.9 (34.3-47.8)	Yes
QNNOTOB3: Percentage of students who did not currently use cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	87.8 (85.1-90.0)	80.4 (78.1-82.5)	70.3 (66.4-73.9)	56.5 (48.8-63.8)	Yes
QNNOTOB2: Percentage of students who did not currently smoke cigarettes or cigars (on at least 1 day during the 30 days before the survey)	89.1 (86.5-91.2)	82.5 (80.3-84.6)	75.1 (71.9-78.1)	61.2 (54.2-67.7)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male**  
**Alcohol/Other Drug Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN42: Percentage of students who drank alcohol before age 13 years (for the first time other than a few sips)	7.3 (5.5-9.5)	9.9 (8.7-11.2)	16.3 (14.0-18.9)	29.5 (23.7-36.1)	Yes
QN43: Percentage of students who currently drank alcohol (at least one drink of alcohol on at least 1 day during the 30 days before the survey)	20.1 (17.4-23.0)	29.5 (26.7-32.4)	33.7 (30.6-36.9)	44.7 (36.6-53.0)	Yes
QN44: Percentage of students who drank five or more drinks of alcohol in a row (within a couple of hours on at least 1 day during the 30 days before the survey)	12.1 (10.0-14.4)	16.9 (14.9-19.1)	20.8 (18.2-23.7)	31.4 (25.1-38.5)	Yes
QN46: Percentage of students who usually obtained the alcohol they drank by someone giving it to them (among students who currently drank alcohol)	44.9 (38.1-51.9)	35.9 (31.4-40.7)	37.6 (31.4-44.3)	29.8 (19.6-42.6)	No
QN48: Percentage of students who tried marijuana before age 13 years (for the first time)	3.3 (2.4-4.4)	5.8 (4.6-7.1)	12.3 (10.4-14.5)	22.1 (16.8-28.6)	Yes
QN49: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	13.3 (11.2-15.8)	22.0 (19.4-24.9)	34.9 (31.3-38.7)	38.7 (32.4-45.5)	Yes
QN50: Percentage of students who ever used cocaine (any form of cocaine, such as powder, crack, or freebase, one or more times during their life)	4.0 (2.8-5.7)	4.2 (3.4-5.1)	9.3 (7.5-11.6)	16.8 (12.5-22.2)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male**  
**Alcohol/Other Drug Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	4.4 (3.2-6.2)	4.9 (4.1-5.8)	7.7 (6.4-9.2)	21.1 (16.1-27.1)	Yes
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China white," one or more times during their life)	2.5 (1.8-3.6)	1.8 (1.3-2.4)	3.0 (2.2-4.1)	10.0 (7.1-13.8)	Yes
QN53: Percentage of students who ever used methamphetamines (also called "speed," "crystal," "crank," or "ice," one or more times during their life)	2.7 (1.7-4.4)	1.9 (1.4-2.5)	3.4 (2.6-4.5)	9.5 (6.8-13.2)	Yes
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	2.7 (2.0-3.7)	3.8 (3.1-4.7)	7.3 (6.0-8.8)	14.3 (10.4-19.4)	Yes
QN55: Percentage of students who ever used synthetic marijuana (also called "K2", "Spice", "fake weed", "King Kong", "Yucatan Fire", "Skunk", or "Moon Rocks", one or more times during their life)	4.8 (3.5-6.6)	8.5 (7.2-10.1)	15.2 (12.9-17.8)	25.3 (20.0-31.5)	Yes
QN57: Percentage of students who ever took prescription drugs without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during their life)	7.5 (6.1-9.2)	12.3 (10.6-14.2)	18.1 (15.6-20.9)	36.8 (31.2-42.7)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Male  
Alcohol/Other Drug Use

Percentage of students who engaged in each risk behavior, by type  
of grades mostly earned in school

Health Risk Behavior	A's	B's	C's	D's/F's	Significant Association*
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN59: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)	13.6 (11.2-16.3)	18.3 (16.3-20.6)	22.7 (19.5-26.3)	35.6 (28.3-43.8)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male  
Sexual Behaviors**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN60: Percentage of students who ever had sexual intercourse	27.4 (24.0-31.0)	39.0 (36.0-42.0)	47.6 (43.9-51.4)	60.5 (53.2-67.3)	Yes
QN61: Percentage of students who had sexual intercourse before age 13 years (for the first time)	2.4 (1.5-3.8)	1.9 (1.4-2.5)	4.0 (3.0-5.2)	10.9 (7.1-16.3)	Yes
QN63: Percentage of students who were currently sexually active (sexual intercourse with at least one person during the 3 months before the survey)	20.6 (17.6-23.9)	29.9 (27.4-32.5)	36.5 (32.9-40.4)	46.0 (38.0-54.2)	Yes
QN64: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	19.4 (14.5-25.3)	15.2 (12.7-18.0)	27.0 (22.6-31.9)	34.8 (24.9-46.3)	Yes
QN65: Percentage of students who used a condom (during last sexual intercourse among students who were currently sexually active)	71.5 (65.9-76.5)	68.4 (63.8-72.7)	66.3 (60.1-72.0)	42.8 (28.8-58.0)	Yes
QN66: Percentage of students who used birth control pills (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	37.6 (30.4-45.4)	29.3 (24.9-34.0)	24.2 (19.3-30.0)	22.1 (12.1-36.8)	No

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male  
Sexual Behaviors**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNIUDIMP: Percentage of students who used an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	2.2 (1.2-4.2)	3.4 (2.4-5.0)	3.1 (1.9-5.1)	1.0 (0.2-4.1)	No
QNSHPARG: Percentage of students who used a shot (e.g., Depo-Provera), patch (e.g., OrthoEvra), or birth control ring (e.g., NuvaRing) (during last sexual intercourse among students who were currently sexually active)	1.7 (0.9-3.5)	2.1 (1.3-3.4)	3.6 (2.1-6.1)	0.8 (0.2-3.1)	No
QNOTHHPL: Percentage of students who used birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	41.6 (34.4-49.2)	34.8 (30.3-39.6)	31.0 (25.7-36.8)	23.8 (13.8-38.0)	No
QNDUALBC: Percentage of students who used both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	24.3 (17.2-33.3)	13.5 (11.1-16.2)	12.7 (8.8-18.1)	2.9 (1.2-6.8)	Yes
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy (during last sexual intercourse among students who were currently sexually active)	4.8 (3.0-7.6)	5.4 (4.0-7.2)	5.6 (3.8-8.4)	21.3 (13.4-32.1)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male**  
**Dietary Behaviors**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNOBESE: Percentage of students who were obese ( $\geq$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart <sup>§</sup> )	9.9 (8.1-12.0)	15.7 (14.0-17.6)	19.3 (16.6-22.2)	24.8 (18.5-32.5)	Yes
QNOWT: Percentage of students who were overweight ( $\geq$ 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart <sup>§</sup> )	14.9 (11.8-18.7)	13.2 (11.7-14.9)	17.5 (14.2-21.4)	14.3 (10.6-18.9)	No
QN70: Percentage of students who were trying to lose weight	21.9 (19.3-24.8)	26.6 (24.0-29.3)	32.2 (28.9-35.7)	35.0 (26.6-44.5)	Yes
QN77: Percentage of students who did not drink a can, bottle, or glass of soda or pop (not including diet soda or diet pop, during the 7 days before the survey)	35.8 (32.4-39.3)	28.1 (25.8-30.5)	24.5 (21.2-28.1)	17.0 (12.4-22.9)	Yes
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	9.4 (7.9-11.2)	16.5 (14.5-18.8)	24.9 (22.0-28.1)	34.3 (28.2-41.0)	Yes
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	5.9 (4.6-7.6)	11.4 (9.7-13.3)	20.5 (17.8-23.6)	30.0 (24.2-36.6)	Yes
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	2.4 (1.8-3.2)	5.8 (4.7-7.3)	11.4 (9.6-13.5)	18.1 (13.6-23.5)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school,  $p < 0.05$ .

† Confidence interval.

§ Based on reference data from the 2000 CDC Growth Charts.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male**  
**Physical Activity**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN80: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	57.8 (53.9-61.7)	55.8 (52.9-58.5)	48.8 (45.2-52.4)	39.8 (32.1-48.0)	Yes
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	9.7 (7.7-12.2)	9.3 (8.1-10.6)	13.9 (11.6-16.7)	23.3 (17.5-30.3)	Yes
QNPA7DAY: Percentage of students who were physically active at least 60 minutes per day on all 7 days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time (during the 7 days before the survey)	30.1 (26.1-34.3)	30.3 (27.8-33.0)	27.5 (24.2-31.0)	20.8 (16.3-26.3)	Yes
QN81: Percentage of students who watched television 3 or more hours per day (on an average school day)	11.2 (9.5-13.2)	18.8 (16.9-20.7)	24.3 (21.3-27.6)	30.7 (25.2-36.9)	Yes
QN82: Percentage of students who played video or computer games or used a computer 3 or more hours per day (for something that was not school work on an average school day)	31.4 (28.0-35.1)	38.8 (36.1-41.6)	46.8 (43.1-50.5)	54.7 (46.2-62.9)	Yes
QN88: Percentage of students who had 8 or more hours of sleep (on an average school night)	36.4 (32.5-40.5)	32.1 (29.7-34.6)	27.2 (24.2-30.5)	21.2 (16.6-26.8)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Male Other	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
	90.0 (87.8-91.7)	84.3 (82.1-86.3)	77.5 (74.6-80.2)	61.7 (54.8-68.1)	Yes
QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work during the 12 months before the survey)					
	20.1 (17.3-23.3)	20.3 (18.4-22.3)	23.7 (20.3-27.4)	26.7 (20.1-34.4)	No
QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma					

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN90: Percentage of students who never or rarely wear a seat belt when driving (among students who drive a car)	5.4 (3.9-7.4)	6.7 (5.5-8.0)	12.7 (10.3-15.5)	29.2 (21.0-39.1)	Yes
QN91: Percentage of students who have experienced an unwanted sexual advance because of another student's drinking (one or more times during the 12 months before the survey)	3.9 (2.5-6.1)	2.6 (2.0-3.3)	4.6 (3.2-6.6)	9.6 (5.8-15.5)	Yes
QN92: Percentage of students who ever saw or heard adults in their home slap, hit, kick, punch, or hurt each other	8.3 (5.5-12.4)	9.6 (8.2-11.2)	13.9 (11.9-16.1)	21.0 (16.3-26.7)	Yes
QN93: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)	7.1 (5.5-9.2)	8.2 (6.5-10.2)	12.0 (10.2-14.0)	23.7 (18.3-30.0)	Yes
QN94: Percentage of students who ever came to school high on marijuana (during their life)	8.2 (6.7-10.0)	14.9 (13.1-16.8)	26.8 (24.0-29.7)	34.6 (28.0-41.9)	Yes
QN95: Percentage of students who currently took a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during the 30 days before the survey)	4.1 (3.0-5.6)	5.6 (4.6-6.8)	8.7 (7.0-10.8)	24.2 (19.0-30.2)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN96: Percentage of students who ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight (during the 30 days before the survey)	22.5 (19.3-26.2)	22.1 (19.5-25.0)	26.5 (23.1-30.1)	28.9 (22.4-36.5)	No
QN97: Percentage of students who will play on their school's sports team one or more seasons (during the school year of the survey)	69.8 (65.6-73.7)	65.3 (62.5-68.1)	50.9 (46.6-55.2)	32.5 (25.8-39.9)	Yes
QN98: Percentage of students who used an indoor tanning device (such as a sunlamp, sunbed, or tanning booth [not including getting a spray-on tan], one or more times during the 12 months before the survey)	2.1 (1.1-3.8)	2.1 (1.6-2.6)	2.9 (2.1-4.0)	6.6 (4.5-9.7)	Yes
QN99: Percentage of students who saw a doctor or nurse (for a check-up or physical exam when they were not sick or injured during the 12 months before the survey)	84.9 (82.2-87.2)	80.7 (78.2-82.9)	76.1 (73.4-78.6)	67.6 (60.4-74.0)	Yes
QN100: Percentage of students who have someone in their family (a parent, brother, or sister) who is currently in the military (air Force, Army, Marines, National Guard, Navy, or Reserves)	11.6 (9.2-14.4)	13.8 (12.1-15.8)	22.2 (18.6-26.3)	23.5 (17.4-30.9)	Yes
QN101: Percentage of students who reported either of their parents or other adults in their family have been in jail or in prison (during the 12 months before the survey)	4.5 (3.3-5.9)	6.8 (5.8-8.0)	12.0 (9.8-14.7)	19.0 (14.6-24.5)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN102: Percentage of students who strongly agree or agree that their parents or other adults in their family have clear rules and consequences for their behavior	81.9 (78.9-84.5)	78.3 (75.6-80.7)	73.0 (70.4-75.5)	55.2 (48.8-61.4)	Yes
QN103: Percentage of students who went hungry most of the time or always because there was not enough food in their home	1.8 (0.9-3.7)	2.3 (1.5-3.3)	2.2 (1.6-3.1)	8.6 (6.0-12.3)	Yes
QN104: Percentage of students who have ever lived with someone who had a problem with alcohol or drugs	17.8 (15.0-20.9)	25.8 (23.7-28.0)	38.4 (34.7-42.2)	50.4 (43.4-57.3)	Yes
QN105: Percentage of students who have talked with at least one of their parents or guardians about the dangers of tobacco, alcohol, or drug use (during the 12 months before the survey)	52.1 (48.3-55.9)	47.9 (45.2-50.6)	43.8 (39.6-48.1)	40.8 (34.1-47.9)	Yes
QN106: Percentage of students who recall hearing, reading, or seeing a message about avoiding alcohol or other illegal drugs (during the 12 months before the survey)	84.6 (81.9-87.0)	81.1 (79.5-82.7)	73.7 (70.6-76.6)	68.6 (62.3-74.2)	Yes
QN107: Percentage of students who would most likely talk with their parent or other adult family member about their feelings (among students who report having felt sad, empty, hopeless, angry, or anxious)	26.3 (23.1-29.8)	20.8 (18.5-23.3)	15.3 (11.8-19.8)	10.3 (7.0-15.0)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN108: Percentage of students who performed organized community service as a non-paid volunteer (such as serving meals to the elderly, picking up litter, helping out at a hospital, or building homes for the poor, one or more times during the 30 days before the survey)	50.1 (45.9-54.4)	38.9 (36.3-41.6)	28.2 (24.2-32.5)	27.3 (19.5-36.8)	Yes
QN109: Percentage of students who participate in activities run by community groups (including activities run by 4-H, Boys and Girls Club, YMCA, sports clubs, or church groups, for one or more hours during an average week when they are in school)	46.0 (41.4-50.8)	34.6 (31.9-37.5)	23.2 (20.2-26.5)	17.2 (12.7-22.9)	Yes
QN110: Percentage of students who think people are at great risk harming themselves if they smoke one or more packs of cigarettes per day (physically or in other ways)	77.8 (74.3-81.0)	70.6 (68.1-73.0)	60.1 (56.3-63.9)	51.4 (44.1-58.7)	Yes
QN111: Percentage of students who think people are at great risk harming themselves if they have one or two drinks of alcohol nearly every day (physically or in other ways)	25.9 (23.0-28.9)	23.5 (21.1-25.9)	20.2 (17.4-23.3)	16.5 (12.0-22.3)	Yes
QN112: Percentage of students who think people are at great risk of harming themselves (physically or in other ways) if they have five or more drinks of alcohol once or twice a week (beer, wine, or liquor)	49.0 (44.9-53.1)	37.9 (34.8-41.1)	30.7 (26.9-34.9)	25.1 (20.0-31.0)	Yes
QN113: Percentage of students who think people are at great risk of harming themselves if they use marijuana once or twice a week (physically or in other ways)	24.1 (21.0-27.4)	16.4 (14.2-18.9)	12.0 (9.8-14.5)	9.1 (5.3-15.2)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN114: Percentage of students who think people are at great risk of harming themselves if they take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	66.7 (62.7-70.5)	61.0 (58.3-63.7)	53.5 (50.0-56.9)	47.7 (39.9-55.5)	Yes
QN115: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to smoke tobacco	72.9 (69.0-76.4)	63.4 (61.1-65.6)	53.0 (49.5-56.6)	38.2 (29.4-47.8)	Yes
QN116: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to have one or two drinks of alcoholic beverage nearly every day (beer, wine or liquor)	60.8 (57.2-64.2)	51.1 (48.6-53.5)	43.7 (39.7-47.8)	34.2 (27.2-42.0)	Yes
QN117: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to smoke marijuana	52.3 (47.9-56.8)	40.3 (37.2-43.5)	30.6 (27.5-34.0)	21.9 (16.7-28.1)	Yes
QN118: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	83.7 (80.6-86.4)	79.4 (77.4-81.3)	75.1 (72.3-77.7)	60.5 (52.0-68.4)	Yes
QN119: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to smoke tobacco	94.4 (92.5-95.9)	89.2 (87.4-90.8)	84.3 (81.0-87.1)	73.8 (67.5-79.2)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Male  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN120: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to have one or two drinks of an alcoholic beverage nearly every day (beer, wine or liquor)	90.9 (88.1-93.1)	87.5 (85.6-89.2)	85.8 (83.7-87.6)	72.5 (66.0-78.1)	Yes
QN121: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to smoke marijuana	90.5 (88.2-92.3)	81.7 (79.3-83.8)	76.4 (73.3-79.3)	65.4 (58.9-71.2)	Yes
QN122: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	96.0 (94.6-97.1)	94.0 (92.7-95.1)	94.0 (92.6-95.2)	85.5 (79.4-90.0)	Yes
QN123: Percentage of students who approve or strongly approve of someone their age having one or two drinks of alcohol nearly every day (beer, wine, or liquor)	5.8 (4.4-7.7)	7.5 (6.4-8.6)	14.8 (12.1-17.9)	20.5 (16.0-26.0)	Yes
QN124: Percentage of students who think it would be very easy for them to get some cigarettes if they wanted to	28.5 (25.4-31.8)	38.9 (35.8-42.1)	45.5 (41.2-49.9)	51.3 (44.9-57.7)	Yes
QN125: Percentage of students who think it would be very easy for them to get some alcohol if they wanted to (beer, wine, or liquor)	32.9 (29.5-36.4)	38.0 (35.3-40.6)	41.9 (38.1-45.9)	44.1 (37.7-50.6)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Male  
Site-Added

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN126: Percentage of students who think it would be very easy for them to get some marijuana if they wanted to	32.7 (29.2-36.4)	42.8 (40.0-45.7)	50.1 (46.3-53.8)	60.6 (54.2-66.5)	Yes
QN127: Percentage of students who think it would be very easy for them to get some prescription drug without a doctor's prescription if they wanted to (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	11.6 (8.9-15.1)	14.7 (12.4-17.3)	20.1 (17.0-23.6)	26.2 (20.8-32.4)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Female Unintentional Injuries/Violence		Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
Health Risk Behavior	A's	B's	C's	D's/F's		
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI		
QN8: Percentage of students who rarely or never wore a bicycle helmet (among students who had ridden a bicycle during the 12 months before the survey)	34.9 (31.3-38.7)	59.7 (55.1-64.1)	74.5 (69.7-78.8)	75.1 (64.4-83.4)	Yes	
QN9: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)	3.2 (1.9-5.2)	6.4 (5.4-7.5)	11.2 (9.1-13.7)	24.3 (18.6-31.0)	Yes	
QN10: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle one or more times during the 30 days before the survey)	12.4 (10.0-15.2)	17.1 (14.5-19.9)	18.0 (15.1-21.2)	37.9 (30.2-46.3)	Yes	
QN11: Percentage of students who drove when drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	3.4 (2.3-4.9)	5.1 (4.0-6.5)	4.1 (2.6-6.3)	22.6 (14.4-33.6)	Yes	
QN12: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	44.6 (41.0-48.2)	48.7 (44.8-52.6)	42.1 (34.6-50.1)	46.3 (35.5-57.4)	No	
QN16: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	2.3 (1.8-3.0)	5.9 (5.0-7.0)	12.3 (9.2-16.3)	26.9 (20.3-34.9)	Yes	

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Female Unintentional Injuries/Violence	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN19: Percentage of students who were injured in a physical fight (one or more times during the 12 months before the survey; injuries had to be treated by a doctor or nurse)	1.0 (0.7-1.5)	2.2 (1.7-2.9)	5.6 (3.6-8.7)	13.5 (9.2-19.4)	Yes
QN20: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	1.2 (0.9-1.7)	3.4 (2.7-4.1)	5.7 (4.2-7.8)	18.2 (12.8-25.3)	Yes
QN21: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	5.1 (3.6-7.3)	10.3 (8.9-11.9)	17.3 (14.2-21.0)	32.3 (24.9-40.7)	Yes
QN22: Percentage of students who experienced physical dating violence (one or more times during the 12 months before the survey, including being hit, slammed into something, or injured with an object or weapon on purpose by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	4.9 (3.5-6.9)	9.4 (8.0-11.1)	16.4 (12.9-20.6)	26.5 (18.9-35.7)	Yes
QN23: Percentage of students who experienced sexual dating violence (one or more times during the 12 months before the survey, including kissing, touching, or being physically forced to have sexual intercourse when they did not want to by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	12.0 (9.1-15.6)	19.0 (16.2-22.3)	22.3 (18.7-26.2)	37.5 (29.0-46.7)	Yes
QN24: Percentage of students who were bullied on school property (during the 12 months before the survey)	18.5 (16.4-20.9)	29.5 (27.1-32.0)	41.4 (35.2-47.7)	51.7 (43.9-59.4)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Female Unintentional Injuries/Violence		Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
Health Risk Behavior	A's	B's	C's	D's/F's		
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI		
QN25: Percentage of students who were electronically bullied (including being bullied through e-mail, chat rooms, instant messaging, websites, or texting during the 12 months before the survey)	16.7 (14.5-19.1)	29.4 (26.9-31.9)	35.8 (30.7-41.1)	47.7 (39.5-56.0)	Yes	
QN26: Percentage of students who felt sad or hopeless (almost every day for 2 or more weeks in a row so that they stopped doing some usual activities during the 12 months before the survey)	25.1 (22.6-27.6)	40.0 (36.1-43.9)	56.0 (50.7-61.0)	72.4 (64.8-79.0)	Yes	
QN27: Percentage of students who seriously considered attempting suicide (during the 12 months before the survey)	12.4 (10.6-14.4)	21.0 (18.9-23.3)	32.6 (28.1-37.5)	53.9 (45.8-61.7)	Yes	
QN29: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	3.9 (3.2-4.7)	8.6 (7.4-10.1)	18.6 (14.8-23.0)	36.3 (28.9-44.4)	Yes	
QN30: Percentage of students who attempted suicide that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	1.0 (0.7-1.5)	2.9 (2.1-4.0)	9.4 (6.4-13.7)	14.9 (9.8-22.0)	Yes	

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Female  
Tobacco Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN32: Percentage of students who smoked a whole cigarette before age 13 years (for the first time)	1.3 (0.9-1.7)	3.5 (2.8-4.2)	7.1 (5.3-9.4)	17.8 (12.4-24.9)	Yes
QN33: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	2.5 (2.0-3.2)	7.9 (6.9-9.1)	17.6 (14.5-21.2)	34.8 (27.4-43.0)	Yes
QNFRCIG: Percentage of students who currently frequently smoked cigarettes (on 20 or more days during the 30 days before the survey)	0.7 (0.4-1.1)	2.7 (2.1-3.4)	5.9 (3.7-9.2)	16.2 (10.7-23.8)	Yes
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	0.5 (0.3-0.8)	1.9 (1.4-2.5)	3.6 (2.5-5.1)	13.3 (8.2-20.8)	Yes
QN34: Percentage of students who smoked more than 10 cigarettes per day (during the 30 days before the survey among students who currently smoked cigarettes on the days they smoked)	7.5 (2.9-17.9)	5.3 (3.0-9.2)	3.1 (1.3-7.1)	23.1 (12.6-38.4)	Yes
QN35: Percentage of students who usually obtained their own cigarettes by buying them in a store or gas station (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	8.9 (3.8-19.7)	7.0 (4.1-12.0)	5.0 (2.2-11.0)	7.7 (2.7-19.9)	No

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Female  
Tobacco Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNCIGINT: Percentage of students who usually obtained their own cigarettes by buying on the internet (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	1.3 (0.2-8.7)	0.8 (0.1-5.2)	1.2 (0.2-8.2)	7.8 (2.2-24.5)	Yes
QN36: Percentage of students who tried to quit smoking cigarettes (among students who currently smoked cigarettes during the 12 months before the survey)	41.7 (30.2-54.2)	40.1 (33.6-47.0)	65.4 (55.9-73.9)	57.5 (42.8-70.9)	Yes
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, or dip on at least 1 day during the 30 days before the survey)	0.7 (0.4-1.1)	1.1 (0.7-1.6)	2.6 (1.7-4.0)	9.0 (5.5-14.3)	Yes
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars on at least 1 day during the 30 days before the survey)	2.6 (2.1-3.3)	6.2 (5.1-7.5)	9.7 (7.1-13.0)	19.4 (14.0-26.2)	Yes
QN40: Percentage of students who currently used electronic vapor products (e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens such as blu, NJOY, or Starbuzz on at least 1 day during the 30 days before the survey)	13.5 (11.4-15.8)	27.3 (24.6-30.2)	35.8 (31.3-40.4)	55.9 (47.7-63.9)	Yes
QNTOB4: Percentage of students who currently used tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	15.0 (12.9-17.5)	30.3 (27.5-33.2)	41.2 (36.4-46.2)	66.5 (58.6-73.6)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Female  
Tobacco Use

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNTOB3: Percentage of students who currently used cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	4.6 (3.8-5.5)	12.2 (10.7-13.9)	22.2 (18.7-26.2)	39.8 (31.7-48.4)	Yes
QNTOB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	4.4 (3.6-5.2)	11.8 (10.4-13.5)	21.2 (17.8-25.0)	39.2 (31.1-47.9)	Yes
QNNOTOB4: Percentage of students who did not currently use tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	85.2 (82.8-87.3)	70.3 (67.3-73.1)	60.0 (55.0-64.8)	35.2 (27.8-43.4)	Yes
QNNOTOB3: Percentage of students who did not currently use cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	95.6 (94.7-96.3)	88.0 (86.3-89.5)	78.5 (74.6-81.9)	61.9 (53.1-70.0)	Yes
QNNOTOB2: Percentage of students who did not currently smoke cigarettes or cigars (on at least 1 day during the 30 days before the survey)	95.8 (94.9-96.5)	88.4 (86.8-89.9)	79.5 (75.7-82.9)	62.5 (53.7-70.6)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Female**  
**Alcohol/Other Drug Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN42: Percentage of students who drank alcohol before age 13 years (for the first time other than a few sips)	4.6 (3.7-5.7)	9.2 (7.2-11.6)	12.7 (10.4-15.3)	30.7 (23.4-39.0)	Yes
QN43: Percentage of students who currently drank alcohol (at least one drink of alcohol on at least 1 day during the 30 days before the survey)	22.5 (20.1-25.2)	36.2 (32.7-39.8)	39.3 (33.7-45.3)	54.1 (45.8-62.2)	Yes
QN44: Percentage of students who drank five or more drinks of alcohol in a row (within a couple of hours on at least 1 day during the 30 days before the survey)	11.0 (9.3-13.1)	18.0 (15.8-20.5)	22.4 (18.1-27.5)	31.8 (24.5-40.1)	Yes
QN46: Percentage of students who usually obtained the alcohol they drank by someone giving it to them (among students who currently drank alcohol)	54.5 (49.0-60.0)	48.0 (44.1-51.9)	38.3 (31.8-45.2)	34.7 (24.4-46.7)	Yes
QN48: Percentage of students who tried marijuana before age 13 years (for the first time)	1.1 (0.8-1.6)	4.4 (2.7-7.0)	7.6 (4.9-11.6)	19.5 (13.8-26.8)	Yes
QN49: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	10.1 (8.7-11.7)	23.4 (20.6-26.4)	33.0 (28.3-38.0)	48.3 (40.2-56.5)	Yes
QN50: Percentage of students who ever used cocaine (any form of cocaine, such as powder, crack, or freebase, one or more times during their life)	1.2 (0.9-1.7)	2.8 (2.0-3.8)	5.2 (3.9-7.0)	18.2 (12.2-26.2)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Female Alcohol/Other Drug Use		Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
Health Risk Behavior	A's	B's	C's	D's/F's		
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI		
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	2.8 (2.2-3.5)	5.7 (4.9-6.7)	9.7 (7.8-12.0)	25.3 (19.0-32.8)	Yes	
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China white," one or more times during their life)	0.6 (0.3-1.1)	0.9 (0.6-1.4)	1.8 (1.1-3.0)	10.9 (6.9-16.9)	Yes	
QN53: Percentage of students who ever used methamphetamines (also called "speed," "crystal," "crank," or "ice," one or more times during their life)	0.8 (0.5-1.1)	1.1 (0.7-1.6)	1.9 (1.2-3.1)	6.7 (3.9-11.2)	Yes	
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	1.2 (0.8-1.6)	3.7 (2.8-5.0)	6.3 (3.8-10.3)	11.9 (7.5-18.2)	Yes	
QN55: Percentage of students who ever used synthetic marijuana (also called "K2", "Spice", "fake weed", "King Kong", "Yucatan Fire", "Skunk", or "Moon Rocks", one or more times during their life)	2.7 (2.1-3.4)	7.8 (6.6-9.2)	13.9 (10.8-17.7)	33.5 (25.7-42.2)	Yes	
QN57: Percentage of students who ever took prescription drugs without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during their life)	7.1 (5.9-8.5)	13.8 (12.2-15.6)	18.6 (15.6-22.0)	34.4 (26.6-43.2)	Yes	

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Female  
Alcohol/Other Drug Use

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN59: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)	9.7 (7.9-11.8)	14.0 (12.0-16.3)	20.4 (16.4-25.1)	31.5 (24.6-39.3)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

#### Female Sexual Behaviors

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN60: Percentage of students who ever had sexual intercourse	27.4 (24.2-30.9)	42.5 (38.9-46.3)	52.0 (46.0-57.9)	67.0 (58.5-74.6)	Yes
QN61: Percentage of students who had sexual intercourse before age 13 years (for the first time)	0.9 (0.6-1.4)	1.5 (1.1-2.1)	2.7 (1.6-4.3)	10.3 (6.3-16.3)	Yes
QN63: Percentage of students who were currently sexually active (sexual intercourse with at least one person during the 3 months before the survey)	22.8 (19.8-26.0)	35.9 (32.7-39.1)	43.0 (37.4-48.7)	52.3 (43.8-60.7)	Yes
QN64: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	10.7 (8.1-13.9)	14.5 (12.0-17.5)	19.2 (15.0-24.1)	36.8 (26.8-48.1)	Yes
QN65: Percentage of students who used a condom (during last sexual intercourse among students who were currently sexually active)	64.6 (59.1-69.7)	53.9 (50.3-57.4)	48.1 (40.6-55.8)	47.8 (37.1-58.7)	Yes
QN66: Percentage of students who used birth control pills (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	51.4 (44.0-58.8)	36.1 (32.0-40.4)	28.7 (23.0-35.2)	26.9 (16.6-40.5)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Female  
Sexual Behaviors**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNIUDIMP: Percentage of students who used an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	4.6 (3.2-6.6)	7.4 (5.4-10.2)	8.7 (6.1-12.3)	8.7 (4.2-16.9)	No
QNSHPARG: Percentage of students who used a shot (e.g., Depo-Provera), patch (e.g., OrthoEvra), or birth control ring (e.g., NuvaRing) (during last sexual intercourse among students who were currently sexually active)	3.6 (2.4-5.5)	6.2 (4.6-8.1)	7.5 (3.4-15.8)	4.3 (1.5-12.0)	No
QNOTHHPL: Percentage of students who used birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	59.7 (52.0-66.9)	49.7 (45.2-54.2)	44.9 (38.0-52.1)	39.9 (28.3-52.8)	No
QNDUALBC: Percentage of students who used both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	33.3 (26.7-40.7)	18.6 (15.7-21.8)	12.3 (9.1-16.4)	17.0 (9.5-28.6)	Yes
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy (during last sexual intercourse among students who were currently sexually active)	3.4 (2.2-5.3)	7.4 (5.8-9.4)	9.3 (6.5-13.2)	13.4 (7.8-22.0)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Female  
Dietary Behaviors**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNOBESE: Percentage of students who were obese ( $\geq$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart <sup>§</sup> )	4.6 (3.8-5.5)	8.7 (7.2-10.3)	13.4 (10.9-16.3)	18.5 (13.4-25.0)	Yes
QNOWT: Percentage of students who were overweight ( $\geq$ 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart <sup>§</sup> )	11.4 (9.9-13.0)	15.4 (13.3-17.9)	16.5 (12.8-21.0)	23.1 (16.8-30.9)	Yes
QN70: Percentage of students who were trying to lose weight	54.6 (50.8-58.3)	65.5 (62.2-68.7)	69.5 (65.4-73.4)	71.3 (63.7-77.9)	Yes
QN77: Percentage of students who did not drink a can, bottle, or glass of soda or pop (not including diet soda or diet pop, during the 7 days before the survey)	55.9 (52.6-59.1)	41.7 (39.1-44.3)	32.9 (28.1-38.0)	32.1 (25.2-39.9)	Yes
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	4.0 (2.9-5.4)	7.4 (6.4-8.5)	13.0 (10.5-16.0)	22.3 (16.4-29.6)	Yes
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	1.5 (1.1-2.0)	4.7 (3.9-5.6)	9.8 (7.7-12.3)	17.0 (11.8-23.8)	Yes
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	0.9 (0.6-1.3)	2.4 (1.9-3.0)	4.5 (3.1-6.3)	10.5 (6.7-16.2)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school,  $p < 0.05$ .

† Confidence interval.

§ Based on reference data from the 2000 CDC Growth Charts.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Female  
Physical Activity

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN80: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	49.2 (45.6-52.8)	36.2 (32.3-40.2)	32.3 (26.3-39.0)	26.6 (20.1-34.3)	Yes
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	11.1 (9.2-13.3)	15.0 (12.9-17.3)	21.4 (18.2-25.1)	28.5 (21.9-36.3)	Yes
QNPA7DAY: Percentage of students who were physically active at least 60 minutes per day on all 7 days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time (during the 7 days before the survey)	15.4 (13.6-17.4)	14.7 (12.5-17.3)	12.6 (9.0-17.3)	14.9 (10.0-21.7)	No
QN81: Percentage of students who watched television 3 or more hours per day (on an average school day)	14.1 (12.0-16.4)	20.9 (18.3-23.8)	24.3 (20.4-28.7)	30.5 (24.1-37.9)	Yes
QN82: Percentage of students who played video or computer games or used a computer 3 or more hours per day (for something that was not school work on an average school day)	31.2 (28.1-34.4)	38.9 (36.3-41.6)	44.3 (39.2-49.5)	48.6 (40.3-57.0)	Yes
QN88: Percentage of students who had 8 or more hours of sleep (on an average school night)	24.5 (22.1-27.1)	26.1 (22.2-30.3)	20.5 (17.4-24.0)	21.5 (15.6-28.7)	No

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Female Other	Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
		A's	B's	C's	D's/F's	
		% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
	QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work during the 12 months before the survey)	90.2 (88.5-91.6)	83.5 (81.6-85.3)	75.8 (72.2-79.1)	67.9 (60.0-74.8)	Yes
	QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma	22.6 (19.5-26.1)	22.6 (20.5-25.0)	23.6 (19.9-27.7)	22.2 (15.9-30.0)	No

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Female  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN90: Percentage of students who never or rarely wear a seat belt when driving (among students who drive a car)	2.1 (1.5-2.9)	5.0 (4.0-6.3)	10.5 (7.4-14.7)	21.5 (14.7-30.2)	Yes
QN91: Percentage of students who have experienced an unwanted sexual advance because of another student's drinking (one or more times during the 12 months before the survey)	5.0 (4.1-6.1)	6.9 (5.7-8.2)	11.7 (8.1-16.6)	17.4 (11.4-25.6)	Yes
QN92: Percentage of students who ever saw or heard adults in their home slap, hit, kick, punch, or hurt each other	12.2 (10.2-14.4)	17.1 (15.4-19.0)	27.6 (21.8-34.3)	42.4 (35.0-50.1)	Yes
QN93: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)	17.2 (15.3-19.3)	29.3 (25.4-33.4)	43.8 (37.8-50.0)	59.6 (51.2-67.4)	Yes
QN94: Percentage of students who ever came to school high on marijuana (during their life)	5.7 (4.2-7.6)	14.5 (12.7-16.6)	23.7 (20.2-27.7)	42.6 (34.7-50.9)	Yes
QN95: Percentage of students who currently took a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during the 30 days before the survey)	2.5 (2.0-3.1)	6.9 (5.7-8.2)	8.6 (6.7-10.9)	21.0 (15.2-28.3)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Female  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN96: Percentage of students who ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight (during the 30 days before the survey)	50.0 (45.9-54.1)	54.2 (51.7-56.7)	62.0 (56.9-66.8)	64.6 (56.8-71.8)	Yes
QN97: Percentage of students who will play on their school's sports team one or more seasons (during the school year of the survey)	66.2 (63.0-69.2)	56.5 (53.4-59.5)	41.5 (36.0-47.3)	28.6 (21.8-36.6)	Yes
QN98: Percentage of students who used an indoor tanning device (such as a sunlamp, sunbed, or tanning booth [not including getting a spray-on tan], one or more times during the 12 months before the survey)	6.2 (4.9-7.9)	11.0 (9.1-13.1)	11.4 (8.3-15.5)	13.3 (8.4-20.5)	Yes
QN99: Percentage of students who saw a doctor or nurse (for a check-up or physical exam when they were not sick or injured during the 12 months before the survey)	86.5 (84.8-88.0)	79.2 (76.5-81.7)	71.6 (66.9-75.9)	68.4 (61.0-75.0)	Yes
QN100: Percentage of students who have someone in their family (a parent, brother, or sister) who is currently in the military (air Force, Army, Marines, National Guard, Navy, or Reserves)	10.0 (8.3-12.0)	13.3 (11.7-15.0)	23.0 (17.0-30.4)	17.1 (12.0-23.9)	Yes
QN101: Percentage of students who reported either of their parents or other adults in their family have been in jail or in prison (during the 12 months before the survey)	5.4 (3.9-7.3)	9.5 (8.3-10.8)	17.0 (14.3-20.1)	30.4 (23.8-37.9)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Female  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN102: Percentage of students who strongly agree or agree that their parents or other adults in their family have clear rules and consequences for their behavior	84.1 (81.8-86.1)	75.9 (73.3-78.3)	69.8 (65.5-73.7)	47.1 (39.0-55.3)	Yes
QN103: Percentage of students who went hungry most of the time or always because there was not enough food in their home	0.9 (0.5-1.3)	1.7 (1.1-2.7)	4.2 (2.3-7.3)	10.5 (6.7-16.0)	Yes
QN104: Percentage of students who have ever lived with someone who had a problem with alcohol or drugs	22.9 (20.3-25.8)	34.6 (31.8-37.6)	50.5 (44.8-56.2)	57.5 (49.4-65.2)	Yes
QN105: Percentage of students who have talked with at least one of their parents or guardians about the dangers of tobacco, alcohol, or drug use (during the 12 months before the survey)	57.1 (53.5-60.6)	52.1 (48.7-55.6)	51.5 (45.9-57.0)	43.6 (35.9-51.7)	Yes
QN106: Percentage of students who recall hearing, reading, or seeing a message about avoiding alcohol or other illegal drugs (during the 12 months before the survey)	86.4 (84.5-88.1)	79.6 (77.1-81.9)	74.8 (71.0-78.3)	71.9 (64.1-78.5)	Yes
QN107: Percentage of students who would most likely talk with their parent or other adult family member about their feelings (among students who report having felt sad, empty, hopeless, angry, or anxious)	26.7 (23.3-30.4)	17.4 (14.8-20.4)	14.9 (11.7-18.8)	7.4 (4.0-13.5)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Female  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN108: Percentage of students who performed organized community service as a non-paid volunteer (such as serving meals to the elderly, picking up litter, helping out at a hospital, or building homes for the poor, one or more times during the 30 days before the survey)	60.4 (57.1-63.6)	43.5 (40.5-46.6)	29.0 (22.9-36.0)	26.4 (19.9-34.1)	Yes
QN109: Percentage of students who participate in activities run by community groups (including activities run by 4-H, Boys and Girls Club, YMCA, sports clubs, or church groups, for one or more hours during an average week when they are in school)	52.3 (48.4-56.2)	36.6 (33.7-39.7)	24.3 (19.9-29.3)	16.5 (11.7-22.8)	Yes
QN110: Percentage of students who think people are at great risk harming themselves if they smoke one or more packs of cigarettes per day (physically or in other ways)	81.7 (79.2-83.9)	71.1 (68.3-73.8)	60.3 (54.0-66.2)	60.2 (52.2-67.6)	Yes
QN111: Percentage of students who think people are at great risk harming themselves if they have one or two drinks of alcohol nearly every day (physically or in other ways)	34.5 (31.1-38.1)	28.8 (26.1-31.7)	29.3 (23.3-36.2)	20.2 (14.3-27.7)	Yes
QN112: Percentage of students who think people are at great risk of harming themselves (physically or in other ways) if they have five or more drinks of alcohol once or twice a week (beer, wine, or liquor)	50.5 (46.8-54.2)	41.3 (38.3-44.4)	33.9 (29.5-38.5)	32.6 (25.9-40.1)	Yes
QN113: Percentage of students who think people are at great risk of harming themselves if they use marijuana once or twice a week (physically or in other ways)	24.8 (22.1-27.8)	18.4 (15.8-21.3)	9.2 (7.3-11.5)	8.3 (4.8-14.0)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Female  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN114: Percentage of students who think people are at great risk of harming themselves if they take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	72.1 (68.4-75.4)	65.9 (63.3-68.4)	56.5 (51.0-61.9)	51.8 (43.9-59.7)	Yes
QN115: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to smoke tobacco	81.8 (79.2-84.2)	69.5 (66.7-72.1)	52.6 (46.9-58.3)	45.3 (36.8-54.1)	Yes
QN116: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to have one or two drinks of alcoholic beverage nearly every day (beer, wine or liquor)	75.6 (73.1-78.0)	61.2 (57.8-64.5)	50.2 (44.5-55.8)	38.3 (31.1-46.1)	Yes
QN117: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to smoke marijuana	58.1 (54.1-62.0)	43.0 (39.6-46.4)	29.4 (23.6-36.0)	21.5 (15.7-28.7)	Yes
QN118: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	91.7 (90.1-93.1)	85.1 (82.6-87.3)	78.1 (74.1-81.6)	64.3 (55.9-72.0)	Yes
QN119: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to smoke tobacco	96.2 (94.6-97.3)	92.9 (91.6-94.0)	88.3 (84.8-91.1)	78.7 (71.1-84.7)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Female  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN120: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to have one or two drinks of an alcoholic beverage nearly every day (beer, wine or liquor)	95.4 (94.3-96.2)	92.8 (91.7-93.8)	89.1 (85.8-91.7)	79.1 (71.4-85.2)	Yes
QN121: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to smoke marijuana	92.3 (91.0-93.4)	83.4 (80.8-85.8)	74.4 (69.3-79.0)	63.5 (55.2-71.0)	Yes
QN122: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	98.0 (97.3-98.5)	97.3 (96.7-97.9)	93.9 (90.7-96.0)	86.8 (80.5-91.2)	Yes
QN123: Percentage of students who approve or strongly approve of someone their age having one or two drinks of alcohol nearly every day (beer, wine, or liquor)	4.6 (3.0-7.1)	6.0 (4.9-7.4)	8.2 (6.4-10.4)	18.2 (12.8-25.3)	Yes
QN124: Percentage of students who think it would be very easy for them to get some cigarettes if they wanted to	25.6 (22.3-29.3)	36.0 (32.7-39.3)	45.0 (39.0-51.2)	60.5 (52.6-67.8)	Yes
QN125: Percentage of students who think it would be very easy for them to get some alcohol if they wanted to (beer, wine, or liquor)	36.0 (32.7-39.6)	39.4 (35.7-43.1)	35.5 (30.5-40.8)	46.8 (39.0-54.9)	No

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Female  
Site-Added

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN126: Percentage of students who think it would be very easy for them to get some marijuana if they wanted to	29.9 (26.8-33.3)	44.3 (41.1-47.6)	51.1 (45.1-57.1)	62.6 (55.1-69.5)	Yes
QN127: Percentage of students who think it would be very easy for them to get some prescription drug without a doctor's prescription if they wanted to (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	9.3 (7.8-11.0)	13.2 (11.7-15.0)	19.9 (15.5-25.2)	30.0 (23.2-37.8)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

White* Unintentional Injuries/Violence	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN8: Percentage of students who rarely or never wore a bicycle helmet (among students who had ridden a bicycle during the 12 months before the survey)	34.8 (32.2-37.6)	59.7 (57.4-62.0)	77.2 (73.7-80.4)	79.6 (72.1-85.5)	Yes
QN9: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)	2.7 (2.1-3.5)	6.0 (5.2-6.8)	11.5 (9.9-13.2)	25.8 (20.8-31.6)	Yes
QN10: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle one or more times during the 30 days before the survey)	12.7 (11.0-14.7)	15.0 (13.3-16.9)	15.6 (13.6-17.8)	30.1 (25.2-35.6)	Yes
QN11: Percentage of students who drove when drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	3.7 (2.8-4.8)	5.5 (4.6-6.5)	6.1 (4.8-7.7)	15.9 (11.2-22.1)	Yes
QN12: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	42.4 (39.3-45.5)	46.1 (43.0-49.2)	41.6 (37.1-46.3)	41.2 (32.8-50.2)	Yes
QN16: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	2.8 (2.0-3.8)	4.3 (3.6-5.2)	7.5 (6.0-9.2)	10.0 (7.2-13.9)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

White* Unintentional Injuries/Violence	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN19: Percentage of students who were injured in a physical fight (one or more times during the 12 months before the survey; injuries had to be treated by a doctor or nurse)	1.7 (1.1-2.5)	2.5 (2.1-3.0)	5.3 (4.1-6.8)	8.4 (6.1-11.6)	Yes
QN20: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	2.5 (1.9-3.4)	4.0 (3.4-4.7)	8.5 (7.2-9.9)	18.1 (13.8-23.5)	Yes
QN21: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	3.0 (2.3-3.9)	5.7 (4.9-6.6)	7.8 (6.3-9.6)	13.9 (9.9-19.2)	Yes
QN22: Percentage of students who experienced physical dating violence (one or more times during the 12 months before the survey, including being hit, slammed into something, or injured with an object or weapon on purpose by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	4.7 (3.5-6.2)	6.7 (5.7-7.9)	10.5 (8.3-13.0)	12.5 (8.9-17.3)	Yes
QN23: Percentage of students who experienced sexual dating violence (one or more times during the 12 months before the survey, including kissing, touching, or being physically forced to have sexual intercourse when they did not want to by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	8.9 (6.9-11.4)	11.8 (10.1-13.7)	11.8 (10.1-13.8)	18.2 (13.2-24.6)	Yes
QN24: Percentage of students who were bullied on school property (during the 12 months before the survey)	15.4 (13.7-17.3)	22.4 (20.8-24.1)	28.6 (25.3-32.2)	34.4 (29.6-39.7)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

White* Unintentional Injuries/Violence	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
<b>Health Risk Behavior</b>					
QN25: Percentage of students who were electronically bullied (including being bullied through e-mail, chat rooms, instant messaging, websites, or texting during the 12 months before the survey)	12.6 (11.1-14.2)	19.7 (18.1-21.4)	22.1 (19.0-25.4)	29.0 (24.0-34.4)	Yes
QN26: Percentage of students who felt sad or hopeless (almost every day for 2 or more weeks in a row so that they stopped doing some usual activities during the 12 months before the survey)	19.0 (17.3-20.9)	27.4 (25.3-29.5)	34.7 (31.8-37.6)	43.9 (38.5-49.3)	Yes
QN27: Percentage of students who seriously considered attempting suicide (during the 12 months before the survey)	9.1 (8.1-10.1)	14.7 (13.4-16.2)	19.9 (17.7-22.3)	35.7 (30.6-41.0)	Yes
QN29: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	2.8 (2.3-3.4)	5.7 (4.8-6.8)	10.4 (8.6-12.5)	20.4 (15.4-26.7)	Yes
QN30: Percentage of students who attempted suicide that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	0.8 (0.5-1.1)	1.8 (1.4-2.5)	4.5 (3.2-6.5)	6.8 (4.4-10.4)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

White* Tobacco Use	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN32: Percentage of students who smoked a whole cigarette before age 13 years (for the first time)	1.8 (1.2-2.5)	3.3 (2.8-3.9)	8.4 (7.0-9.9)	17.0 (13.2-21.6)	Yes
QN33: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	3.0 (2.4-3.6)	7.5 (6.6-8.5)	16.1 (14.2-18.3)	30.7 (25.1-37.0)	Yes
QNFRICIG: Percentage of students who currently frequently smoked cigarettes (on 20 or more days during the 30 days before the survey)	0.8 (0.6-1.2)	2.6 (2.1-3.1)	6.5 (5.2-8.2)	13.2 (9.6-17.9)	Yes
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	0.5 (0.4-0.8)	1.7 (1.4-2.2)	4.1 (3.2-5.2)	11.2 (7.8-15.9)	Yes
QN34: Percentage of students who smoked more than 10 cigarettes per day (during the 30 days before the survey among students who currently smoked cigarettes on the days they smoked)	8.2 (4.1-15.6)	5.9 (3.8-9.0)	7.7 (4.9-11.7)	19.4 (12.7-28.6)	Yes
QN35: Percentage of students who usually obtained their own cigarettes by buying them in a store or gas station (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	13.9 (7.8-23.5)	9.1 (6.2-13.1)	14.1 (10.0-19.6)	19.3 (12.3-28.9)	No

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

White\*  
Tobacco Use

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
QNCIGINT: Percentage of students who usually obtained their own cigarettes by buying on the internet (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	0.0	1.7 (0.6-4.4)	1.2 (0.3-4.9)	2.4 (0.6-9.7)	No
QN36: Percentage of students who tried to quit smoking cigarettes (among students who currently smoked cigarettes during the 12 months before the survey)	40.1 (31.6-49.2)	43.2 (37.7-48.9)	55.9 (49.6-62.1)	48.0 (37.2-58.9)	Yes
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, or dip on at least 1 day during the 30 days before the survey)	2.1 (1.5-3.0)	5.0 (4.2-5.9)	9.7 (7.9-11.8)	18.1 (12.7-25.0)	Yes
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars on at least 1 day during the 30 days before the survey)	5.8 (4.7-7.1)	11.0 (9.6-12.6)	14.8 (12.6-17.3)	23.3 (18.4-29.0)	Yes
QN40: Percentage of students who currently used electronic vapor products (e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens such as blu, NJOY, or Starbuzz on at least 1 day during the 30 days before the survey)	13.2 (11.6-14.9)	26.0 (24.1-28.0)	36.3 (32.8-40.0)	52.8 (46.3-59.1)	Yes
QNTOB4: Percentage of students who currently used tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	16.7 (14.7-18.9)	31.2 (29.0-33.5)	43.8 (39.9-47.8)	62.8 (56.7-68.5)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

White* Tobacco Use	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QNTOB3: Percentage of students who currently used cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	7.9 (6.8-9.3)	16.6 (14.9-18.4)	28.1 (24.9-31.5)	42.2 (35.4-49.3)	Yes
QNTOB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	7.3 (6.2-8.7)	15.2 (13.6-17.0)	24.0 (21.3-26.8)	37.8 (31.7-44.4)	Yes
QNNOTOB4: Percentage of students who did not currently use tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	83.9 (81.7-85.9)	69.6 (67.3-71.8)	57.3 (53.2-61.2)	38.7 (32.7-45.0)	Yes
QNNOTOB3: Percentage of students who did not currently use cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	92.6 (91.3-93.7)	84.1 (82.3-85.8)	72.7 (69.2-75.9)	59.3 (52.1-66.1)	Yes
QNNOTOB2: Percentage of students who did not currently smoke cigarettes or cigars (on at least 1 day during the 30 days before the survey)	93.2 (91.9-94.2)	85.5 (83.7-87.1)	76.6 (73.8-79.3)	63.3 (56.7-69.5)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

White\*  
Alcohol/Other Drug Use

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
QN42: Percentage of students who drank alcohol before age 13 years (for the first time other than a few sips)	5.0 (3.9-6.4)	8.7 (7.6-10.0)	14.2 (12.3-16.3)	27.1 (21.9-33.1)	Yes
QN43: Percentage of students who currently drank alcohol (at least one drink of alcohol on at least 1 day during the 30 days before the survey)	21.8 (19.7-24.1)	33.1 (30.2-36.2)	35.8 (32.6-39.2)	45.5 (37.7-53.4)	Yes
QN44: Percentage of students who drank five or more drinks of alcohol in a row (within a couple of hours on at least 1 day during the 30 days before the survey)	11.4 (9.8-13.3)	17.4 (15.6-19.4)	20.9 (18.3-23.7)	29.6 (23.6-36.4)	Yes
QN46: Percentage of students who usually obtained the alcohol they drank by someone giving it to them (among students who currently drank alcohol)	52.7 (48.8-56.6)	42.7 (39.7-45.8)	39.7 (34.3-45.2)	33.7 (23.8-45.4)	Yes
QN48: Percentage of students who tried marijuana before age 13 years (for the first time)	1.8 (1.3-2.3)	4.7 (3.6-6.0)	10.2 (8.5-12.2)	16.8 (12.9-21.6)	Yes
QN49: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	11.1 (9.7-12.6)	22.4 (20.1-24.9)	34.0 (30.6-37.7)	38.7 (32.6-45.1)	Yes
QN50: Percentage of students who ever used cocaine (any form of cocaine, such as powder, crack, or freebase, one or more times during their life)	1.9 (1.4-2.7)	3.3 (2.7-4.0)	7.8 (6.3-9.6)	13.3 (9.8-17.8)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

White* Alcohol/Other Drug Use	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	3.0 (2.3-3.8)	5.0 (4.4-5.7)	8.3 (7.1-9.6)	19.4 (14.8-25.1)	Yes
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China white," one or more times during their life)	1.0 (0.6-1.7)	1.2 (0.9-1.6)	2.4 (1.7-3.2)	7.0 (4.7-10.1)	Yes
QN53: Percentage of students who ever used methamphetamines (also called "speed," "crystal," "crank," or "ice," one or more times during their life)	1.2 (0.7-1.9)	1.3 (1.0-1.7)	2.6 (2.0-3.5)	4.9 (3.3-7.5)	Yes
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	1.5 (1.1-2.0)	3.4 (2.8-4.3)	6.9 (5.5-8.7)	9.3 (6.6-12.9)	Yes
QN55: Percentage of students who ever used synthetic marijuana (also called "K2", "Spice", "fake weed", "King Kong", "Yucatan Fire", "Skunk", or "Moon Rocks", one or more times during their life)	3.2 (2.5-4.2)	8.0 (6.9-9.2)	14.6 (12.5-17.1)	24.1 (19.2-29.8)	Yes
QN57: Percentage of students who ever took prescription drugs without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during their life)	6.7 (5.7-7.8)	12.8 (11.3-14.3)	18.2 (15.9-20.7)	34.5 (29.0-40.5)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

White* Alcohol/Other Drug Use	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN59: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)	10.8 (9.4-12.3)	16.2 (14.7-17.9)	21.7 (18.7-25.1)	33.1 (26.2-40.8)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

White* Sexual Behaviors	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN60: Percentage of students who ever had sexual intercourse	27.5 (24.7-30.5)	40.9 (38.0-43.9)	49.0 (45.4-52.6)	59.8 (53.1-66.1)	Yes
QN61: Percentage of students who had sexual intercourse before age 13 years (for the first time)	1.2 (0.8-1.9)	1.4 (1.1-1.8)	3.4 (2.5-4.5)	5.8 (3.8-8.6)	Yes
QN63: Percentage of students who were currently sexually active (sexual intercourse with at least one person during the 3 months before the survey)	22.1 (19.4-25.0)	32.8 (30.4-35.4)	38.9 (35.6-42.3)	43.3 (36.2-50.7)	Yes
QN64: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	12.8 (10.0-16.3)	14.3 (12.4-16.5)	23.1 (19.4-27.3)	32.7 (23.6-43.3)	Yes
QN65: Percentage of students who used a condom (during last sexual intercourse among students who were currently sexually active)	68.0 (63.4-72.2)	61.0 (58.1-63.9)	58.5 (53.0-63.9)	41.2 (30.5-52.8)	Yes
QN66: Percentage of students who used birth control pills (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	46.9 (41.6-52.3)	33.0 (29.8-36.3)	25.3 (21.0-30.2)	28.5 (19.0-40.4)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

White* Sexual Behaviors	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QNIUDIMP: Percentage of students who used an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	3.8 (2.7-5.3)	5.8 (4.5-7.5)	5.7 (4.2-7.8)	4.0 (1.9-8.3)	Yes
QNSHPARG: Percentage of students who used a shot (e.g., Depo-Provera), patch (e.g., OrthoEvra), or birth control ring (e.g., NuvaRing) (during last sexual intercourse among students who were currently sexually active)	3.0 (2.1-4.4)	4.4 (3.4-5.7)	5.6 (3.2-9.7)	0.7 (0.1-4.8)	Yes
QNOTHHPL: Percentage of students who used birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	53.8 (48.4-59.0)	43.2 (39.9-46.7)	36.6 (31.5-42.1)	33.3 (23.8-44.2)	Yes
QNDUALBC: Percentage of students who used both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	31.0 (25.7-36.9)	16.1 (14.1-18.4)	11.5 (8.5-15.4)	8.0 (4.3-14.4)	Yes
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy (during last sexual intercourse among students who were currently sexually active)	3.5 (2.4-5.1)	5.8 (4.7-7.2)	7.4 (5.6-9.7)	19.4 (12.3-29.4)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

White* Dietary Behaviors	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QNOBESE: Percentage of students who were obese (>= 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart¶)	6.7 (5.7-7.8)	12.3 (11.2-13.6)	17.4 (15.4-19.6)	22.5 (16.8-29.4)	Yes
QNOWT: Percentage of students who were overweight (>= 85th percentile but <95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart¶)	12.7 (11.1-14.6)	14.3 (12.7-16.1)	17.5 (14.1-21.3)	17.1 (13.2-21.9)	Yes
QN70: Percentage of students who were trying to lose weight	41.6 (38.7-44.6)	45.4 (43.4-47.4)	46.3 (43.1-49.6)	49.4 (41.3-57.5)	Yes
QN77: Percentage of students who did not drink a can, bottle, or glass of soda or pop (not including diet soda or diet pop, during the 7 days before the survey)	47.9 (45.1-50.7)	34.8 (32.8-36.9)	27.5 (24.7-30.6)	21.8 (17.0-27.5)	Yes
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	6.0 (5.0-7.3)	12.2 (10.8-13.9)	20.5 (18.1-23.1)	29.4 (23.9-35.5)	Yes
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	3.1 (2.5-3.9)	8.2 (7.1-9.4)	16.4 (14.2-18.9)	25.1 (19.8-31.2)	Yes
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	1.3 (1.0-1.7)	4.0 (3.3-4.9)	8.4 (7.1-9.9)	14.6 (10.9-19.4)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

¶Based on reference data from the 2000 CDC Growth Charts.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

White* Physical Activity	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN80: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	53.7 (51.0-56.4)	46.6 (43.5-49.6)	42.7 (39.3-46.1)	35.0 (28.1-42.6)	Yes
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	9.3 (8.0-10.7)	11.5 (10.2-12.9)	16.1 (13.9-18.5)	25.5 (20.1-31.7)	Yes
QNPA7DAY: Percentage of students who were physically active at least 60 minutes per day on all 7 days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	21.2 (19.1-23.4)	23.1 (21.1-25.2)	21.2 (18.6-24.2)	18.0 (14.0-22.8)	Yes
QN81: Percentage of students who watched television 3 or more hours per day (on an average school day)	12.5 (10.9-14.2)	19.4 (17.9-21.0)	23.9 (20.9-27.1)	28.8 (23.8-34.4)	Yes
QN82: Percentage of students who played video or computer games or used a computer 3 or more hours per day (for something that was not school work on an average school day)	30.5 (28.1-33.0)	38.7 (36.7-40.8)	45.8 (42.9-48.7)	55.1 (47.3-62.6)	Yes
QN88: Percentage of students who had 8 or more hours of sleep (on an average school night)	29.9 (27.6-32.3)	29.4 (27.0-32.0)	24.8 (22.3-27.6)	22.8 (18.4-27.8)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

White* Other	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school					Significant Association†
	A's	B's	C's	D's/F's		
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI		
<b>Health Risk Behavior</b>						
QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work during the 12 months before the survey)	90.9 (89.6-92.1)	84.9 (83.1-86.4)	78.9 (76.4-81.2)	66.6 (60.5-72.1)		Yes
QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma	21.6 (19.5-23.9)	21.2 (19.9-22.6)	23.2 (20.5-26.3)	20.9 (16.3-26.3)		No

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

White\*  
Site-Added

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association <sup>†</sup>
	A's	B's	C's	D's/F's	
	% 95% CI <sup>‡</sup>	% 95% CI	% 95% CI	% 95% CI	
QN90: Percentage of students who never or rarely wear a seat belt when driving (among students who drive a car)	2.7 (2.0-3.6)	5.4 (4.6-6.4)	11.5 (9.3-14.0)	24.1 (16.9-33.2)	Yes
QN91: Percentage of students who have experienced an unwanted sexual advance because of another student's drinking (one or more times during the 12 months before the survey)	4.3 (3.5-5.4)	4.5 (3.8-5.4)	7.0 (5.2-9.3)	9.2 (5.5-14.9)	Yes
QN92: Percentage of students who ever saw or heard adults in their home slap, hit, kick, punch, or hurt each other	9.5 (7.8-11.5)	12.5 (11.2-13.8)	18.6 (15.7-21.9)	23.9 (19.2-29.4)	Yes
QN93: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)	12.8 (11.4-14.4)	18.7 (16.7-21.0)	24.1 (20.9-27.7)	32.9 (27.2-39.1)	Yes
QN94: Percentage of students who ever came to school high on marijuana (during their life)	5.8 (5.0-6.7)	14.5 (13.0-16.1)	25.8 (23.3-28.5)	33.3 (27.4-39.8)	Yes
QN95: Percentage of students who currently took a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during the 30 days before the survey)	2.7 (2.2-3.4)	5.8 (5.0-6.8)	8.1 (6.7-9.8)	20.4 (15.7-26.2)	Yes

\*Non-Hispanic.

<sup>†</sup>Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

<sup>‡</sup>Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**White\*  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association <sup>†</sup>
	A's	B's	C's	D's/F's	
	% 95% CI <sup>‡</sup>	% 95% CI	% 95% CI	% 95% CI	
QN96: Percentage of students who ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight (during the 30 days before the survey)	39.0 (35.4-42.7)	38.2 (36.2-40.2)	39.9 (36.5-43.4)	41.4 (34.5-48.5)	Yes
QN97: Percentage of students who will play on their school's sports team one or more seasons (during the school year of the survey)	68.5 (66.2-70.7)	61.7 (58.9-64.4)	46.7 (42.3-51.1)	28.0 (22.1-34.7)	Yes
QN98: Percentage of students who used an indoor tanning device (such as a sunlamp, sunbed, or tanning booth [not including getting a spray-on tan], one or more times during the 12 months before the survey)	4.5 (3.6-5.8)	6.4 (5.4-7.6)	5.9 (4.5-7.7)	5.5 (3.5-8.6)	Yes
QN99: Percentage of students who saw a doctor or nurse (for a check-up or physical exam when they were not sick or injured during the 12 months before the survey)	86.6 (85.1-88.0)	80.7 (78.8-82.4)	75.3 (72.5-77.9)	70.4 (64.5-75.7)	Yes
QN100: Percentage of students who have someone in their family (a parent, brother, or sister) who is currently in the military (air Force, Army, Marines, National Guard, Navy, or Reserves)	10.0 (8.7-11.6)	13.3 (12.0-14.7)	22.0 (18.4-26.2)	17.6 (13.7-22.3)	Yes
QN101: Percentage of students who reported either of their parents or other adults in their family have been in jail or in prison (during the 12 months before the survey)	4.1 (3.5-4.9)	7.5 (6.6-8.6)	13.1 (11.2-15.4)	19.1 (15.1-23.9)	Yes

\*Non-Hispanic.

<sup>†</sup>Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

<sup>‡</sup>Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

White\*  
Site-Added

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association <sup>†</sup>
	A's	B's	C's	D's/F's	
	% 95% CI <sup>‡</sup>	% 95% CI	% 95% CI	% 95% CI	
QN102: Percentage of students who strongly agree or agree that their parents or other adults in their family have clear rules and consequences for their behavior	83.3 (81.3-85.0)	77.5 (75.5-79.4)	72.0 (69.3-74.6)	51.8 (46.5-57.0)	Yes
QN103: Percentage of students who went hungry most of the time or always because there was not enough food in their home	1.0 (0.6-1.9)	1.9 (1.4-2.7)	2.2 (1.6-2.9)	6.4 (4.2-9.5)	Yes
QN104: Percentage of students who have ever lived with someone who had a problem with alcohol or drugs	20.6 (18.5-22.9)	30.3 (28.2-32.4)	43.9 (39.9-47.9)	52.4 (46.5-58.2)	Yes
QN105: Percentage of students who have talked with at least one of their parents or guardians about the dangers of tobacco, alcohol, or drug use (during the 12 months before the survey)	55.7 (52.9-58.5)	50.7 (48.0-53.4)	46.0 (42.0-50.0)	42.8 (36.3-49.5)	Yes
QN106: Percentage of students who recall hearing, reading, or seeing a message about avoiding alcohol or other illegal drugs (during the 12 months before the survey)	86.5 (84.9-87.9)	81.3 (79.6-82.9)	74.8 (72.2-77.2)	70.8 (65.2-75.8)	Yes
QN107: Percentage of students who would most likely talk with their parent or other adult family member about their feelings (among students who report having felt sad, empty, hopeless, angry, or anxious)	26.9 (24.1-29.9)	19.2 (17.4-21.1)	15.9 (12.8-19.6)	9.5 (6.5-13.6)	Yes

\*Non-Hispanic.

<sup>†</sup>Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

<sup>‡</sup>Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

White\*  
Site-Added

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
QN108: Percentage of students who performed organized community service as a non-paid volunteer (such as serving meals to the elderly, picking up litter, helping out at a hospital, or building homes for the poor, one or more times during the 30 days before the survey)	57.1 (54.5-59.7)	41.4 (38.9-43.8)	28.6 (25.3-32.1)	28.7 (21.7-37.0)	Yes
QN109: Percentage of students who participate in activities run by community groups (including activities run by 4-H, Boys and Girls Club, YMCA, sports clubs, or church groups, for one or more hours during an average week when they are in school)	49.8 (46.2-53.5)	35.7 (33.4-38.1)	22.7 (20.1-25.6)	17.0 (12.8-22.2)	Yes
QN110: Percentage of students who think people are at great risk harming themselves if they smoke one or more packs of cigarettes per day (physically or in other ways)	81.3 (79.3-83.1)	71.9 (70.0-73.7)	62.1 (58.3-65.8)	58.2 (51.7-64.3)	Yes
QN111: Percentage of students who think people are at great risk harming themselves if they have one or two drinks of alcohol nearly every day (physically or in other ways)	30.4 (28.0-32.9)	25.5 (23.9-27.2)	24.1 (20.3-28.3)	17.6 (13.2-23.1)	Yes
QN112: Percentage of students who think people are at great risk of harming themselves (physically or in other ways) if they have five or more drinks of alcohol once or twice a week (beer, wine, or liquor)	49.8 (46.9-52.8)	39.5 (37.7-41.4)	32.2 (29.3-35.2)	28.2 (23.5-33.4)	Yes
QN113: Percentage of students who think people are at great risk of harming themselves if they use marijuana once or twice a week (physically or in other ways)	23.9 (21.5-26.4)	16.9 (14.8-19.2)	11.0 (9.3-12.9)	9.7 (6.0-15.3)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**White\*  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association <sup>†</sup>
	A's	B's	C's	D's/F's	
	% 95% CI <sup>‡</sup>	% 95% CI	% 95% CI	% 95% CI	
QN114: Percentage of students who think people are at great risk of harming themselves if they take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	70.7 (67.9-73.3)	64.2 (61.9-66.5)	55.1 (51.6-58.5)	52.3 (45.3-59.2)	Yes
QN115: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to smoke tobacco	78.7 (76.1-81.2)	65.9 (63.9-67.9)	53.5 (50.1-56.9)	41.5 (33.4-50.1)	Yes
QN116: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to have one or two drinks of alcoholic beverage nearly every day (beer, wine or liquor)	70.0 (67.7-72.3)	56.0 (53.5-58.5)	46.9 (42.5-51.3)	36.0 (29.7-42.9)	Yes
QN117: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to smoke marijuana	55.5 (51.4-59.5)	41.4 (38.4-44.4)	30.6 (26.9-34.5)	21.6 (16.9-27.3)	Yes
QN118: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	89.5 (87.9-90.9)	83.0 (81.3-84.5)	77.3 (74.7-79.6)	63.5 (55.9-70.5)	Yes
QN119: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to smoke tobacco	95.7 (94.4-96.7)	91.2 (89.9-92.3)	86.0 (83.4-88.3)	77.5 (71.8-82.3)	Yes

\*Non-Hispanic.

<sup>†</sup>Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

<sup>‡</sup>Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**White\*  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
QN120: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to have one or two drinks of an alcoholic beverage nearly every day (beer, wine or liquor)	94.1 (92.7-95.2)	90.4 (89.1-91.6)	87.7 (85.9-89.2)	78.0 (72.2-83.0)	Yes
QN121: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to smoke marijuana	91.9 (90.7-93.0)	82.7 (80.9-84.4)	75.9 (73.2-78.5)	68.1 (62.1-73.6)	Yes
QN122: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	97.8 (97.1-98.4)	96.0 (95.3-96.7)	94.6 (93.3-95.6)	89.6 (83.9-93.4)	Yes
QN123: Percentage of students who approve or strongly approve of someone their age having one or two drinks of alcohol nearly every day (beer, wine, or liquor)	4.2 (2.9-5.9)	6.1 (5.3-7.0)	12.0 (10.0-14.3)	15.9 (12.2-20.5)	Yes
QN124: Percentage of students who think it would be very easy for them to get some cigarettes if they wanted to	26.7 (23.9-29.6)	37.3 (34.7-39.9)	45.1 (41.0-49.2)	51.1 (45.9-56.3)	Yes
QN125: Percentage of students who think it would be very easy for them to get some alcohol if they wanted to (beer, wine, or liquor)	35.7 (33.2-38.3)	38.7 (36.5-41.0)	39.8 (36.3-43.4)	42.5 (37.3-47.8)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

White\*  
Site-Added

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
QN126: Percentage of students who think it would be very easy for them to get some marijuana if they wanted to	31.5 (29.2-34.0)	43.6 (41.4-45.8)	50.0 (46.2-53.7)	60.0 (54.6-65.2)	Yes
QN127: Percentage of students who think it would be very easy for them to get some prescription drug without a doctor's prescription if they wanted to (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	10.1 (8.6-11.9)	13.6 (12.1-15.4)	19.6 (16.6-23.0)	23.4 (18.8-28.7)	Yes

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

<b>Black*</b> <b>Unintentional Injuries/Violence</b>  <b>Health Risk Behavior</b>	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				<b>Significant Association†</b>
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
QN8: Percentage of students who rarely or never wore a bicycle helmet (among students who had ridden a bicycle during the 12 months before the survey)	-	62.9 (49.1-74.9)	92.2 (80.4-97.2)	-	
QN9: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)	4.6 (1.4-13.7)	12.0 (6.6-21.0)	16.9 (7.2-34.7)	-	
QN10: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle one or more times during the 30 days before the survey)	8.2 (3.0-20.6)	11.5 (6.9-18.5)	22.0 (11.0-39.0)	-	
QN11: Percentage of students who drove when drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	-	4.1 (1.0-15.3)	-	-	
QN12: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	-	52.6 (37.6-67.2)	-	-	
QN16: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	7.1 (1.5-27.3)	5.7 (2.4-12.7)	11.5 (4.8-25.2)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
QN19: Percentage of students who were injured in a physical fight (one or more times during the 12 months before the survey; injuries had to be treated by a doctor or nurse)	3.2 (0.4-19.6)	6.0 (3.0-11.9)	9.2 (3.1-24.4)	-	
QN20: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	9.5 (2.8-27.8)	12.8 (7.4-21.2)	22.5 (11.7-38.8)	-	
QN21: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	11.0 (3.2-31.3)	7.9 (3.7-15.9)	0.5 (0.1-3.3)	-	
QN22: Percentage of students who experienced physical dating violence (one or more times during the 12 months before the survey, including being hit, slammed into something, or injured with an object or weapon on purpose by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	-	8.1 (3.2-19.3)	3.3 (0.4-20.3)	-	
QN23: Percentage of students who experienced sexual dating violence (one or more times during the 12 months before the survey, including kissing, touching, or being physically forced to have sexual intercourse when they did not want to by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	-	10.4 (4.7-21.2)	15.4 (5.8-35.0)	-	
QN24: Percentage of students who were bullied on school property (during the 12 months before the survey)	11.9 (4.3-28.7)	16.2 (9.8-25.5)	9.5 (3.8-21.5)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

<b>Black*</b> <b>Unintentional Injuries/Violence</b>  <b>Health Risk Behavior</b>	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				<b>Significant Association†</b>
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
QN25: Percentage of students who were electronically bullied (including being bullied through e-mail, chat rooms, instant messaging, websites, or texting during the 12 months before the survey)	9.7 (3.8-22.9)	14.4 (8.2-24.2)	7.8 (2.8-19.7)	-	
QN26: Percentage of students who felt sad or hopeless (almost every day for 2 or more weeks in a row so that they stopped doing some usual activities during the 12 months before the survey)	23.1 (11.6-40.7)	24.3 (16.6-34.2)	15.0 (7.1-29.0)	-	
QN27: Percentage of students who seriously considered attempting suicide (during the 12 months before the survey)	10.8 (4.0-26.2)	12.4 (6.8-21.5)	8.3 (3.3-19.3)	-	
QN29: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	3.3 (1.0-10.1)	4.4 (1.7-10.7)	5.2 (1.5-16.2)	-	
QN30: Percentage of students who attempted suicide that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	1.2 (0.2-8.0)	1.6 (0.3-7.5)	2.2 (0.3-14.5)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Black* Tobacco Use	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
<b>Health Risk Behavior</b>					
QN32: Percentage of students who smoked a whole cigarette before age 13 years (for the first time)	5.4 (1.1-23.4)	5.3 (2.0-13.4)	5.2 (1.6-15.3)	-	
QN33: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	2.4 (0.6-9.4)	7.1 (3.4-14.0)	7.5 (2.5-20.3)	-	
QNFRICIG: Percentage of students who currently frequently smoked cigarettes (on 20 or more days during the 30 days before the survey)	0.0	2.0 (0.5-8.3)	0.7 (0.1-5.1)	-	
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	0.0	2.0 (0.5-8.3)	0.0	-	
QN34: Percentage of students who smoked more than 10 cigarettes per day (during the 30 days before the survey among students who currently smoked cigarettes on the days they smoked)	-	-	-	-	
QN35: Percentage of students who usually obtained their own cigarettes by buying them in a store or gas station (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	-	-	-	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Tobacco Use	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN36: Percentage of students who tried to quit smoking cigarettes (among students who currently smoked cigarettes during the 12 months before the survey)	-	-	-	-	
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, or dip on at least 1 day during the 30 days before the survey)	8.0 (1.9-27.3)	5.8 (2.5-13.1)	6.6 (2.3-17.5)	-	
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars on at least 1 day during the 30 days before the survey)	6.7 (1.3-28.0)	7.9 (4.0-15.3)	15.2 (7.7-27.9)	-	
QN40: Percentage of students who currently used electronic vapor products (e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens such as blu, NJOY, or Starbuzz on at least 1 day during the 30 days before the survey)	19.3 (9.6-34.8)	26.5 (17.5-38.1)	33.2 (16.0-56.5)	-	
QNTOB4: Percentage of students who currently used tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	25.7 (13.6-43.3)	31.3 (21.5-43.0)	37.6 (20.6-58.4)	-	
QNTOB3: Percentage of students who currently used cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	10.2 (3.3-27.6)	11.7 (6.7-19.8)	15.7 (8.0-28.3)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Tobacco Use	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QNTOB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	7.8 (1.8-27.5)	10.3 (5.6-18.2)	15.7 (8.0-28.3)	-	
QNNOTOB4: Percentage of students who did not currently use tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	79.4 (63.4-89.5)	69.9 (58.1-79.6)	63.9 (42.2-81.2)	-	
QNNOTOB3: Percentage of students who did not currently use cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	96.0 (87.7-98.8)	89.2 (81.4-94.0)	86.4 (73.9-93.5)	-	
QNNOTOB2: Percentage of students who did not currently smoke cigarettes or cigars (on at least 1 day during the 30 days before the survey)	97.6 (90.6-99.4)	89.7 (81.8-94.4)	86.4 (73.9-93.5)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Alcohol/Other Drug Use	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN42: Percentage of students who drank alcohol before age 13 years (for the first time other than a few sips)	12.5 (4.9-28.6)	13.0 (7.5-21.6)	23.3 (12.7-38.8)	-	
QN43: Percentage of students who currently drank alcohol (at least one drink of alcohol on at least 1 day during the 30 days before the survey)	15.4 (6.9-30.7)	27.4 (19.0-37.7)	25.8 (14.8-41.0)	-	
QN44: Percentage of students who drank five or more drinks of alcohol in a row (within a couple of hours on at least 1 day during the 30 days before the survey)	12.9 (5.5-27.3)	13.1 (7.9-21.0)	15.8 (8.3-28.1)	-	
QN46: Percentage of students who usually obtained the alcohol they drank by someone giving it to them (among students who currently drank alcohol)	-	38.5 (21.2-59.4)	-	-	
QN48: Percentage of students who tried marijuana before age 13 years (for the first time)	4.4 (0.9-18.9)	5.9 (2.8-12.3)	7.4 (2.4-20.4)	-	
QN49: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	15.1 (6.9-30.1)	27.7 (18.7-38.9)	37.7 (23.2-54.8)	-	
QN50: Percentage of students who ever used cocaine (any form of cocaine, such as powder, crack, or freebase, one or more times during their life)	10.7 (3.3-29.4)	4.1 (1.7-9.6)	6.2 (2.1-17.0)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Alcohol/Other Drug Use	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	15.1 (5.9-33.6)	2.9 (1.3-6.5)	10.0 (3.8-24.1)	-	
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China white," one or more times during their life)	7.6 (1.8-27.2)	0.7 (0.1-4.6)	4.7 (1.2-16.3)	-	
QN53: Percentage of students who ever used methamphetamines (also called "speed," "crystal," "crank," or "ice," one or more times during their life)	8.3 (2.2-26.9)	2.3 (0.6-7.9)	4.7 (1.2-16.3)	-	
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	2.2 (0.5-8.7)	3.1 (1.2-7.5)	5.5 (1.7-16.8)	-	
QN55: Percentage of students who ever used synthetic marijuana (also called "K2", "Spice", "fake weed", "King Kong", "Yucatan Fire", "Skunk", or "Moon Rocks", one or more times during their life)	9.2 (2.6-27.7)	10.7 (5.9-18.7)	13.5 (5.6-29.2)	-	
QN57: Percentage of students who ever took prescription drugs without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during their life)	16.2 (6.6-34.6)	14.9 (9.0-23.5)	7.7 (2.9-18.9)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Alcohol/Other Drug Use	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN59: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)	16.6 (7.1-34.1)	13.2 (7.4-22.4)	19.3 (9.5-35.2)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Sexual Behaviors	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN60: Percentage of students who ever had sexual intercourse	30.5 (17.1-48.3)	30.7 (20.4-43.5)	41.3 (24.5-60.3)	-	
QN61: Percentage of students who had sexual intercourse before age 13 years (for the first time)	10.2 (3.1-29.2)	2.0 (0.6-6.7)	0.0	-	
QN63: Percentage of students who were currently sexually active (sexual intercourse with at least one person during the 3 months before the survey)	18.8 (9.2-34.5)	24.8 (17.1-34.5)	29.7 (16.0-48.3)	-	
QN64: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	-	29.6 (14.7-50.6)	-	-	
QN65: Percentage of students who used a condom (during last sexual intercourse among students who were currently sexually active)	-	49.6 (30.3-69.1)	-	-	
QN66: Percentage of students who used birth control pills (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	-	33.2 (16.4-55.7)	-	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Sexual Behaviors	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QNIUDIMP: Percentage of students who used an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	-	3.3 (0.5-20.2)	-	-	
QNSHPARG: Percentage of students who used a shot (e.g., Depo-Provera), patch (e.g., OrthoEvra), or birth control ring (e.g., NuvaRing) (during last sexual intercourse among students who were currently sexually active)	-	11.0 (2.8-34.4)	-	-	
QNOTHHPL: Percentage of students who used birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	-	47.5 (28.4-67.3)	-	-	
QNDUALBC: Percentage of students who used both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	-	17.0 (5.6-41.1)	-	-	
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy (during last sexual intercourse among students who were currently sexually active)	-	7.1 (1.6-26.0)	-	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Dietary Behaviors	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QNOBESE: Percentage of students who were obese ( $\geq$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart¶)	5.9 (1.1-26.1)	9.3 (4.4-18.8)	12.0 (5.1-25.7)	-	
QNOWT: Percentage of students who were overweight ( $\geq$ 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart¶)	14.3 (5.4-32.9)	16.1 (9.7-25.7)	23.5 (7.4-54.2)	-	
QN70: Percentage of students who were trying to lose weight	24.2 (10.8-45.8)	48.7 (37.9-59.6)	31.7 (15.3-54.6)	-	
QN77: Percentage of students who did not drink a can, bottle, or glass of soda or pop (not including diet soda or diet pop, during the 7 days before the survey)	51.7 (36.0-67.1)	42.8 (33.5-52.6)	31.4 (14.8-54.8)	-	
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	11.2 (3.8-28.5)	11.2 (6.4-18.8)	15.6 (7.3-30.1)	-	
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	9.7 (2.9-28.1)	7.9 (4.0-14.8)	15.6 (7.3-30.1)	-	
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	5.6 (1.1-24.1)	7.1 (3.4-14.1)	10.1 (3.8-24.0)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school,  $p < 0.05$ .

‡Confidence interval.

¶Based on reference data from the 2000 CDC Growth Charts.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Physical Activity	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN80: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	32.5 (18.6-50.5)	35.6 (24.2-48.9)	58.4 (39.1-75.5)	-	
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	19.3 (8.6-37.6)	24.4 (16.0-35.4)	18.5 (9.0-34.2)	-	
QNPA7DAY: Percentage of students who were physically active at least 60 minutes per day on all 7 days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time (during the 7 days before the survey)	16.4 (7.3-32.8)	22.9 (13.1-37.0)	23.0 (12.0-39.6)	-	
QN81: Percentage of students who watched television 3 or more hours per day (on an average school day)	25.6 (13.1-44.0)	26.1 (17.9-36.4)	38.4 (20.9-59.4)	-	
QN82: Percentage of students who played video or computer games or used a computer 3 or more hours per day (for something that was not school work on an average school day)	54.9 (37.2-71.5)	40.9 (30.1-52.6)	46.6 (27.5-66.8)	-	
QN88: Percentage of students who had 8 or more hours of sleep (on an average school night)	27.2 (14.1-46.1)	28.9 (20.2-39.4)	36.8 (22.2-54.3)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association <sup>†</sup>
	A's % 95% CI <sup>§</sup>	B's % 95% CI	C's % 95% CI	D's/F's % 95% CI	
QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work during the 12 months before the survey)	79.5 (60.4-90.8)	72.9 (58.6-83.7)	43.8 (27.3-61.7)	-	
QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma	16.5 (6.9-34.7)	19.6 (12.8-28.8)	20.1 (10.4-35.4)	-	

\*Non-Hispanic.

<sup>†</sup>Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

<sup>§</sup>Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Site-Added	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN90: Percentage of students who never or rarely wear a seat belt when driving (among students who drive a car)	-	7.5 (3.1-16.8)	20.7 (9.4-39.8)	-	
QN91: Percentage of students who have experienced an unwanted sexual advance because of another student's drinking (one or more times during the 12 months before the survey)	7.8 (1.9-27.6)	5.8 (2.1-14.7)	6.4 (1.8-20.1)	-	
QN92: Percentage of students who ever saw or heard adults in their home slap, hit, kick, punch, or hurt each other	32.4 (18.8-49.8)	21.5 (13.3-32.7)	13.1 (6.0-26.2)	-	
QN93: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)	15.1 (6.2-32.2)	10.3 (5.6-18.2)	14.8 (7.2-27.9)	-	
QN94: Percentage of students who ever came to school high on marijuana (during their life)	25.0 (12.7-43.3)	14.7 (8.7-23.8)	24.2 (13.2-40.0)	-	
QN95: Percentage of students who currently took a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during the 30 days before the survey)	1.7 (0.4-6.8)	7.7 (4.0-14.1)	7.2 (2.6-18.6)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Black\*  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
QN96: Percentage of students who ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight (during the 30 days before the survey)	39.8 (22.9-59.6)	36.7 (25.5-49.4)	30.4 (13.3-55.4)	-	
QN97: Percentage of students who will play on their school's sports team one or more seasons (during the school year of the survey)	50.0 (34.3-65.7)	53.6 (42.2-64.6)	78.2 (62.2-88.7)	-	
QN98: Percentage of students who used an indoor tanning device (such as a sunlamp, sunbed, or tanning booth [not including getting a spray-on tan], one or more times during the 12 months before the survey)	3.6 (0.5-21.8)	1.9 (0.6-5.9)	11.9 (5.0-25.5)	-	
QN99: Percentage of students who saw a doctor or nurse (for a check-up or physical exam when they were not sick or injured during the 12 months before the survey)	78.5 (59.8-90.0)	77.2 (65.8-85.7)	61.1 (42.6-76.9)	-	
QN100: Percentage of students who have someone in their family (a parent, brother, or sister) who is currently in the military (air Force, Army, Marines, National Guard, Navy, or Reserves)	13.1 (4.7-31.7)	15.6 (9.2-25.1)	17.4 (9.3-30.3)	-	
QN101: Percentage of students who reported either of their parents or other adults in their family have been in jail or in prison (during the 12 months before the survey)	15.0 (5.7-34.0)	12.3 (7.1-20.4)	18.1 (8.9-33.4)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Site-Added	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN102: Percentage of students who strongly agree or agree that their parents or other adults in their family have clear rules and consequences for their behavior	92.6 (78.7-97.7)	75.1 (63.7-83.9)	73.6 (56.9-85.4)	-	
QN103: Percentage of students who went hungry most of the time or always because there was not enough food in their home	0.0	2.1 (0.6-6.8)	9.1 (3.2-23.2)	-	
QN104: Percentage of students who have ever lived with someone who had a problem with alcohol or drugs	19.4 (8.6-38.1)	21.9 (14.8-31.1)	12.6 (6.5-23.1)	-	
QN105: Percentage of students who have talked with at least one of their parents or guardians about the dangers of tobacco, alcohol, or drug use (during the 12 months before the survey)	35.1 (20.3-53.4)	44.4 (34.4-55.0)	39.0 (21.4-60.0)	-	
QN106: Percentage of students who recall hearing, reading, or seeing a message about avoiding alcohol or other illegal drugs (during the 12 months before the survey)	78.9 (62.5-89.4)	71.3 (60.0-80.4)	58.9 (41.1-74.6)	-	
QN107: Percentage of students who would most likely talk with their parent or other adult family member about their feelings (among students who report having felt sad, empty, hopeless, angry, or anxious)	19.4 (8.6-38.1)	13.4 (7.0-24.3)	17.1 (6.7-37.2)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Black\*  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
QN108: Percentage of students who performed organized community service as a non-paid volunteer (such as serving meals to the elderly, picking up litter, helping out at a hospital, or building homes for the poor, one or more times during the 30 days before the survey)	39.3 (25.2-55.4)	38.7 (28.8-49.7)	18.9 (9.2-35.0)	-	
QN109: Percentage of students who participate in activities run by community groups (including activities run by 4-H, Boys and Girls Club, YMCA, sports clubs, or church groups, for one or more hours during an average week when they are in school)	40.8 (25.1-58.6)	41.5 (30.8-53.1)	46.9 (27.4-67.4)	-	
QN110: Percentage of students who think people are at great risk harming themselves if they smoke one or more packs of cigarettes per day (physically or in other ways)	71.1 (54.1-83.7)	64.9 (54.4-74.2)	48.5 (30.3-67.2)	-	
QN111: Percentage of students who think people are at great risk harming themselves if they have one or two drinks of alcohol nearly every day (physically or in other ways)	53.0 (37.2-68.2)	28.0 (19.0-39.1)	25.2 (12.9-43.3)	-	
QN112: Percentage of students who think people are at great risk of harming themselves (physically or in other ways) if they have five or more drinks of alcohol once or twice a week (beer, wine, or liquor)	60.9 (44.3-75.3)	38.6 (28.7-49.6)	36.8 (21.8-54.8)	-	
QN113: Percentage of students who think people are at great risk of harming themselves if they use marijuana once or twice a week (physically or in other ways)	34.5 (19.4-53.6)	22.2 (13.7-33.9)	6.7 (1.3-28.3)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Site-Added	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN114: Percentage of students who think people are at great risk of harming themselves if they take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	62.1 (45.5-76.3)	54.8 (43.8-65.3)	59.4 (41.6-74.9)	-	
QN115: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to smoke tobacco	68.7 (52.2-81.6)	63.3 (52.9-72.5)	59.7 (37.9-78.3)	-	
QN116: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to have one or two drinks of alcoholic beverage nearly every day (beer, wine or liquor)	59.7 (40.7-76.3)	49.8 (39.5-60.1)	49.9 (30.8-69.1)	-	
QN117: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to smoke marijuana	56.0 (37.5-73.0)	37.2 (27.4-48.2)	28.4 (16.4-44.5)	-	
QN118: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	75.1 (56.4-87.5)	69.3 (59.4-77.7)	73.4 (58.1-84.6)	-	
QN119: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to smoke tobacco	93.3 (75.0-98.5)	82.7 (73.0-89.4)	90.2 (79.2-95.7)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Site-Added	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN120: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to have one or two drinks of an alcoholic beverage nearly every day (beer, wine or liquor)	86.9 (68.8-95.2)	91.0 (82.2-95.7)	86.3 (73.7-93.4)	-	
QN121: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to smoke marijuana	88.2 (70.3-95.9)	82.6 (72.5-89.5)	79.1 (63.1-89.3)	-	
QN122: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	84.9 (65.3-94.4)	92.9 (86.3-96.5)	86.9 (73.9-94.0)	-	
QN123: Percentage of students who approve or strongly approve of someone their age having one or two drinks of alcohol nearly every day (beer, wine, or liquor)	14.4 (5.3-33.5)	11.2 (6.3-19.1)	18.5 (8.4-36.0)	-	
QN124: Percentage of students who think it would be very easy for them to get some cigarettes if they wanted to	30.2 (17.1-47.5)	34.3 (24.0-46.4)	39.2 (21.2-60.7)	-	
QN125: Percentage of students who think it would be very easy for them to get some alcohol if they wanted to (beer, wine, or liquor)	20.2 (9.9-36.8)	34.7 (25.8-44.8)	35.2 (20.7-53.2)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

Black* Site-Added	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association†
	A's	B's	C's	D's/F's	
	% 95% CI‡	% 95% CI	% 95% CI	% 95% CI	
Health Risk Behavior					
QN126: Percentage of students who think it would be very easy for them to get some marijuana if they wanted to	32.9 (20.0-49.0)	46.6 (34.1-59.4)	48.8 (30.9-67.1)	-	
QN127: Percentage of students who think it would be very easy for them to get some prescription drug without a doctor's prescription if they wanted to (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	17.3 (8.0-33.5)	8.4 (4.0-16.9)	29.6 (16.4-47.4)	-	

\*Non-Hispanic.

†Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

‡Confidence interval.

- = Fewer than 30 students in this subgroup.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Hispanic/Latino  
Unintentional Injuries/Violence

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN8: Percentage of students who rarely or never wore a bicycle helmet (among students who had ridden a bicycle during the 12 months before the survey)	64.4 (53.8-73.7)	68.1 (58.3-76.5)	84.2 (74.2-90.8)	87.5 (73.1-94.7)	Yes
QN9: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)	22.7 (7.5-51.7)	8.7 (5.9-12.7)	14.2 (9.0-21.7)	42.7 (31.2-54.9)	Yes
QN10: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle one or more times during the 30 days before the survey)	16.3 (10.4-24.7)	21.3 (16.1-27.6)	19.5 (14.0-26.5)	46.8 (33.9-60.2)	Yes
QN11: Percentage of students who drove when drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	8.5 (4.1-16.8)	8.7 (4.9-15.1)	6.6 (3.1-13.7)	62.8 (49.0-74.7)	Yes
QN12: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	44.3 (32.7-56.6)	41.4 (31.9-51.6)	38.4 (26.7-51.6)	65.5 (50.7-77.7)	Yes
QN16: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	4.9 (2.4-9.9)	4.9 (2.9-8.0)	7.6 (4.0-13.9)	26.0 (14.4-42.2)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

#### Hispanic/Latino

#### Unintentional Injuries/Violence

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN19: Percentage of students who were injured in a physical fight (one or more times during the 12 months before the survey; injuries had to be treated by a doctor or nurse)	8.9 (5.0-15.3)	6.0 (3.8-9.5)	6.5 (3.8-11.0)	28.7 (16.5-45.1)	Yes
QN20: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	6.8 (3.7-12.3)	12.5 (9.0-17.1)	9.8 (6.0-15.7)	38.3 (22.9-56.5)	Yes
QN21: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	24.5 (9.1-51.4)	9.6 (6.3-14.3)	14.1 (9.4-20.8)	24.7 (13.8-40.1)	Yes
QN22: Percentage of students who experienced physical dating violence (one or more times during the 12 months before the survey, including being hit, slammed into something, or injured with an object or weapon on purpose by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	12.3 (7.0-20.6)	10.2 (6.5-15.7)	12.1 (7.0-20.0)	28.4 (14.4-48.2)	No
QN23: Percentage of students who experienced sexual dating violence (one or more times during the 12 months before the survey, including kissing, touching, or being physically forced to have sexual intercourse when they did not want to by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	13.8 (8.5-21.5)	14.4 (9.7-20.9)	14.5 (8.6-23.3)	25.9 (13.0-45.0)	No
QN24: Percentage of students who were bullied on school property (during the 12 months before the survey)	28.7 (12.3-53.5)	17.5 (13.1-23.0)	29.3 (19.4-41.7)	36.0 (21.8-53.2)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

#### Hispanic/Latino

#### Unintentional Injuries/Violence

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN25: Percentage of students who were electronically bullied (including being bullied through e-mail, chat rooms, instant messaging, websites, or texting during the 12 months before the survey)	24.8 (9.2-51.5)	15.0 (10.8-20.5)	23.6 (14.5-36.0)	31.5 (18.6-48.1)	Yes
QN26: Percentage of students who felt sad or hopeless (almost every day for 2 or more weeks in a row so that they stopped doing some usual activities during the 12 months before the survey)	37.7 (21.2-57.6)	30.3 (24.7-36.5)	44.2 (33.7-55.3)	51.9 (31.4-71.8)	Yes
QN27: Percentage of students who seriously considered attempting suicide (during the 12 months before the survey)	24.7 (8.9-52.6)	15.4 (11.4-20.6)	29.7 (19.3-42.7)	32.2 (19.1-48.9)	Yes
QN29: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	5.0 (2.4-9.9)	6.7 (4.2-10.7)	14.7 (9.4-22.2)	35.3 (18.5-56.7)	Yes
QN30: Percentage of students who attempted suicide that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	2.3 (0.8-6.7)	1.7 (0.7-4.3)	6.2 (3.1-11.8)	26.9 (12.2-49.4)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Hispanic/Latino  
Tobacco Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN32: Percentage of students who smoked a whole cigarette before age 13 years (for the first time)	8.2 (4.3-15.0)	9.3 (6.3-13.5)	10.1 (6.2-16.0)	51.3 (37.7-64.7)	Yes
QN33: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	6.6 (3.3-12.7)	9.5 (6.5-13.7)	11.2 (7.2-17.0)	37.7 (22.4-55.8)	Yes
QNFRCIG: Percentage of students who currently frequently smoked cigarettes (on 20 or more days during the 30 days before the survey)	3.6 (1.6-7.8)	4.0 (2.2-7.3)	5.7 (3.0-10.6)	29.9 (17.0-47.0)	Yes
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	3.2 (1.3-7.5)	4.0 (2.2-7.3)	5.1 (2.5-9.9)	23.0 (12.1-39.2)	Yes
QN34: Percentage of students who smoked more than 10 cigarettes per day (during the 30 days before the survey among students who currently smoked cigarettes on the days they smoked)	-	27.8 (13.2-49.2)	-	56.2 (36.9-73.7)	
QN35: Percentage of students who usually obtained their own cigarettes by buying them in a store or gas station (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	-	-	-	-	

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Hispanic/Latino  
Tobacco Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNCIGINT: Percentage of students who usually obtained their own cigarettes by buying on the internet (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	-	-	-	-	
QN36: Percentage of students who tried to quit smoking cigarettes (among students who currently smoked cigarettes during the 12 months before the survey)	-	-	-	31.7 (17.4-50.6)	
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, or dip on at least 1 day during the 30 days before the survey)	6.0 (3.0-11.5)	5.2 (3.2-8.4)	4.9 (2.6-9.3)	37.2 (25.7-50.3)	Yes
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars on at least 1 day during the 30 days before the survey)	7.8 (4.4-13.6)	9.6 (6.5-14.1)	21.8 (13.4-33.4)	39.5 (27.6-52.7)	Yes
QN40: Percentage of students who currently used electronic vapor products (e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens such as blu, NJOY, or Starbuzz on at least 1 day during the 30 days before the survey)	16.0 (9.9-24.7)	30.0 (24.5-36.2)	35.6 (26.4-45.9)	48.3 (35.0-61.9)	Yes
QNTOB4: Percentage of students who currently used tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	19.0 (12.1-28.6)	34.3 (28.1-41.0)	46.3 (35.8-57.1)	60.0 (44.5-73.7)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

- = Fewer than 30 students in this subgroup.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Hispanic/Latino  
Tobacco Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNTOB3: Percentage of students who currently used cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	10.9 (6.4-18.2)	15.2 (11.3-20.3)	27.9 (18.8-39.3)	51.1 (37.6-64.5)	Yes
QNTOB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	10.4 (6.0-17.5)	14.6 (10.8-19.6)	26.5 (17.5-38.0)	49.5 (36.3-62.7)	Yes
QNNOTOB4: Percentage of students who did not currently use tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	82.9 (73.6-89.4)	67.5 (60.8-73.6)	54.7 (43.7-65.3)	41.4 (27.6-56.8)	Yes
QNNOTOB3: Percentage of students who did not currently use cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	90.7 (84.0-94.8)	86.2 (81.3-90.0)	72.4 (60.9-81.5)	50.1 (36.9-63.3)	Yes
QNNOTOB2: Percentage of students who did not currently smoke cigarettes or cigars (on at least 1 day during the 30 days before the survey)	91.2 (84.6-95.2)	86.7 (82.0-90.4)	73.7 (62.1-82.7)	51.8 (38.7-64.6)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Hispanic/Latino  
Alcohol/Other Drug Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN42: Percentage of students who drank alcohol before age 13 years (for the first time other than a few sips)	15.0 (9.3-23.4)	17.9 (13.2-23.7)	19.4 (12.9-28.1)	48.5 (35.0-62.2)	Yes
QN43: Percentage of students who currently drank alcohol (at least one drink of alcohol on at least 1 day during the 30 days before the survey)	26.7 (17.9-37.9)	33.9 (27.5-40.9)	34.4 (24.4-46.0)	53.1 (38.7-67.1)	Yes
QN44: Percentage of students who drank five or more drinks of alcohol in a row (within a couple of hours on at least 1 day during the 30 days before the survey)	16.9 (10.7-25.7)	18.8 (13.7-25.3)	23.0 (14.8-33.9)	41.9 (29.8-55.1)	Yes
QN46: Percentage of students who usually obtained the alcohol they drank by someone giving it to them (among students who currently drank alcohol)	30.3 (18.9-44.7)	33.5 (23.8-44.7)	31.0 (19.3-45.9)	22.2 (11.2-39.1)	No
QN48: Percentage of students who tried marijuana before age 13 years (for the first time)	4.8 (2.3-9.8)	11.0 (7.6-15.7)	12.0 (7.8-18.2)	47.1 (33.5-61.1)	Yes
QN49: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	18.1 (11.6-27.2)	27.9 (21.9-34.7)	33.5 (24.3-44.3)	50.6 (37.0-64.2)	Yes
QN50: Percentage of students who ever used cocaine (any form of cocaine, such as powder, crack, or freebase, one or more times during their life)	6.9 (3.8-12.2)	8.1 (5.4-12.1)	9.5 (5.8-15.2)	36.2 (24.1-50.3)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Hispanic/Latino**  
**Alcohol/Other Drug Use**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	9.3 (5.3-15.9)	7.7 (5.0-11.7)	8.1 (4.7-13.5)	39.0 (26.7-52.9)	Yes
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China white," one or more times during their life)	4.9 (2.5-9.4)	5.6 (3.4-9.2)	3.5 (1.6-7.5)	25.6 (14.0-42.2)	Yes
QN53: Percentage of students who ever used methamphetamines (also called "speed," "crystal," "crank," or "ice," one or more times during their life)	5.5 (2.7-10.7)	4.4 (2.5-7.8)	3.6 (1.7-7.5)	26.9 (14.6-44.2)	Yes
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	6.3 (3.3-11.8)	8.6 (5.7-12.8)	8.7 (5.1-14.3)	36.0 (23.9-50.2)	Yes
QN55: Percentage of students who ever used synthetic marijuana (also called "K2", "Spice", "fake weed", "King Kong", "Yucatan Fire", "Skunk", or "Moon Rocks", one or more times during their life)	9.8 (5.3-17.3)	11.5 (8.0-16.3)	16.2 (9.2-26.8)	43.9 (31.2-57.5)	Yes
QN57: Percentage of students who ever took prescription drugs without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during their life)	16.2 (10.0-25.2)	17.0 (13.0-22.0)	20.8 (13.0-31.6)	50.5 (37.2-63.8)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Hispanic/Latino  
Alcohol/Other Drug Use

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN59: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)	26.7 (10.9-51.9)	17.3 (13.1-22.4)	26.4 (18.0-36.8)	41.1 (29.3-54.1)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

#### Hispanic/Latino Sexual Behaviors

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN60: Percentage of students who ever had sexual intercourse	32.9 (24.7-42.2)	47.6 (39.7-55.7)	57.4 (45.8-68.3)	77.9 (62.2-88.3)	Yes
QN61: Percentage of students who had sexual intercourse before age 13 years (for the first time)	5.8 (2.9-11.1)	5.6 (3.2-9.7)	5.8 (2.5-13.0)	35.4 (18.0-57.8)	Yes
QN63: Percentage of students who were currently sexually active (sexual intercourse with at least one person during the 3 months before the survey)	28.8 (21.2-37.8)	36.7 (29.8-44.3)	42.1 (31.4-53.6)	75.6 (59.4-86.8)	Yes
QN64: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	25.6 (14.2-41.7)	20.1 (13.4-29.2)	30.2 (18.3-45.4)	38.6 (18.8-63.1)	No
QN65: Percentage of students who used a condom (during last sexual intercourse among students who were currently sexually active)	55.5 (38.6-71.1)	60.6 (49.8-70.5)	58.9 (43.6-72.7)	62.2 (38.1-81.5)	No
QN66: Percentage of students who used birth control pills (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	46.0 (30.8-62.0)	22.6 (15.9-31.1)	29.6 (14.5-51.1)	1.3 (0.3-5.6)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

#### Hispanic/Latino Sexual Behaviors

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNIUDIMP: Percentage of students who used an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	3.1 (0.4-19.1)	3.8 (1.2-11.1)	4.5 (1.6-12.0)	2.5 (0.7-8.7)	No
QNSHPARG: Percentage of students who used a shot (e.g., Depo-Provera), patch (e.g., OrthoEvra), or birth control ring (e.g., NuvaRing) (during last sexual intercourse among students who were currently sexually active)	5.1 (1.0-22.2)	3.1 (1.3-7.3)	1.8 (0.4-7.2)	3.5 (0.8-14.5)	No
QNOTHHPL: Percentage of students who used birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	54.2 (37.8-69.7)	29.5 (21.6-38.9)	35.9 (20.0-55.6)	7.3 (2.7-18.3)	Yes
QNDUALBC: Percentage of students who used both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	24.1 (12.1-42.4)	15.3 (9.5-23.6)	18.8 (6.4-43.7)	4.0 (1.2-12.1)	No
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy (during last sexual intercourse among students who were currently sexually active)	8.8 (3.2-21.7)	15.9 (9.7-25.1)	6.4 (2.2-17.0)	19.8 (9.0-38.3)	No

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

#### Hispanic/Latino Dietary Behaviors

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QNOBESE: Percentage of students who were obese ( $\geq$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart <sup>§</sup> )	3.9 (1.9-7.9)	14.3 (10.3-19.5)	20.7 (14.3-28.9)	34.1 (21.0-50.1)	Yes
QNOWT: Percentage of students who were overweight ( $\geq$ 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart <sup>§</sup> )	11.3 (6.7-18.5)	16.3 (12.2-21.4)	11.8 (7.6-17.9)	11.4 (4.7-25.1)	No
QN70: Percentage of students who were trying to lose weight	50.8 (35.5-66.1)	45.6 (38.9-52.4)	47.6 (37.2-58.2)	26.1 (15.2-40.9)	No
QN77: Percentage of students who did not drink a can, bottle, or glass of soda or pop (not including diet soda or diet pop, during the 7 days before the survey)	51.8 (36.7-66.6)	29.5 (24.2-35.3)	27.0 (18.6-37.4)	18.9 (10.2-32.2)	Yes
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	8.9 (5.0-15.5)	12.2 (9.0-16.3)	23.5 (15.6-34.0)	39.6 (27.0-53.7)	Yes
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	5.9 (3.1-10.9)	7.7 (5.2-11.2)	19.3 (11.8-30.0)	34.4 (22.8-48.2)	Yes
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	2.6 (1.2-5.7)	3.7 (2.1-6.2)	12.6 (6.3-23.6)	20.9 (11.1-36.0)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school,  $p < 0.05$ .

† Confidence interval.

§ Based on reference data from the 2000 CDC Growth Charts.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Hispanic/Latino  
Physical Activity**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN80: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	51.1 (35.4-66.5)	44.8 (37.9-51.9)	36.4 (26.1-48.3)	40.3 (28.5-53.4)	Yes
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	26.8 (11.0-52.0)	18.5 (14.1-24.0)	21.6 (15.2-29.8)	23.4 (12.9-38.6)	No
QNPA7DAY: Percentage of students who were physically active at least 60 minutes per day on all 7 days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time (during the 7 days before the survey)	25.3 (17.0-35.8)	18.5 (14.3-23.6)	21.7 (12.7-34.5)	18.9 (10.0-32.7)	No
QN81: Percentage of students who watched television 3 or more hours per day (on an average school day)	19.2 (12.3-28.8)	20.6 (16.0-26.3)	23.5 (16.0-33.3)	38.8 (26.5-52.7)	No
QN82: Percentage of students who played video or computer games or used a computer 3 or more hours per day (for something that was not school work on an average school day)	32.8 (22.1-45.6)	42.2 (35.9-48.7)	54.8 (45.6-63.7)	51.4 (37.9-64.8)	Yes
QN88: Percentage of students who had 8 or more hours of sleep (on an average school night)	19.1 (12.4-28.3)	29.6 (23.4-36.6)	16.7 (11.3-24.0)	11.9 (5.9-22.4)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Hispanic/Latino  
Other

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work during the 12 months before the survey)	84.4 (76.3-90.0)	75.3 (69.4-80.4)	68.1 (57.3-77.3)	44.3 (32.2-57.0)	Yes
QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma	32.7 (16.0-55.3)	24.4 (19.2-30.5)	32.6 (22.1-45.1)	54.2 (35.0-72.3)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Hispanic/Latino  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN90: Percentage of students who never or rarely wear a seat belt when driving (among students who drive a car)	9.6 (4.8-18.0)	9.7 (6.0-15.4)	14.5 (9.0-22.6)	40.9 (27.8-55.4)	Yes
QN91: Percentage of students who have experienced an unwanted sexual advance because of another student's drinking (one or more times during the 12 months before the survey)	6.9 (3.9-11.9)	5.2 (3.3-8.1)	6.9 (3.9-11.9)	24.9 (13.0-42.2)	Yes
QN92: Percentage of students who ever saw or heard adults in their home slap, hit, kick, punch, or hurt each other	30.9 (14.4-54.3)	18.2 (14.0-23.2)	20.6 (14.8-27.9)	44.0 (31.1-57.7)	Yes
QN93: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)	17.8 (11.4-26.8)	14.7 (10.8-19.8)	22.6 (16.2-30.5)	36.5 (21.8-54.2)	Yes
QN94: Percentage of students who ever came to school high on marijuana (during their life)	24.5 (8.9-52.0)	19.7 (15.1-25.3)	27.0 (18.6-37.4)	55.8 (42.2-68.7)	Yes
QN95: Percentage of students who currently took a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during the 30 days before the survey)	9.3 (5.2-16.1)	11.5 (8.2-15.9)	12.5 (6.2-23.6)	38.9 (26.7-52.6)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Hispanic/Latino  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN96: Percentage of students who ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight (during the 30 days before the survey)	51.0 (35.3-66.6)	33.2 (27.3-39.7)	37.5 (27.2-49.1)	31.7 (18.8-48.1)	No
QN97: Percentage of students who will play on their school's sports team one or more seasons (during the school year of the survey)	55.2 (37.5-71.6)	55.9 (48.8-62.8)	47.1 (36.0-58.6)	37.9 (26.0-51.5)	Yes
QN98: Percentage of students who used an indoor tanning device (such as a sunlamp, sunbed, or tanning booth [not including getting a spray-on tan], one or more times during the 12 months before the survey)	6.3 (3.2-12.0)	7.0 (4.6-10.5)	5.2 (2.6-10.2)	20.4 (11.2-34.3)	Yes
QN99: Percentage of students who saw a doctor or nurse (for a check-up or physical exam when they were not sick or injured during the 12 months before the survey)	81.8 (72.8-88.3)	73.5 (67.4-78.8)	71.3 (62.0-79.1)	56.6 (36.2-74.9)	No
QN100: Percentage of students who have someone in their family (a parent, brother, or sister) who is currently in the military (air Force, Army, Marines, National Guard, Navy, or Reserves)	30.1 (13.7-53.9)	17.9 (13.7-23.1)	23.3 (13.8-36.7)	40.2 (20.2-64.0)	No
QN101: Percentage of students who reported either of their parents or other adults in their family have been in jail or in prison (during the 12 months before the survey)	24.6 (8.9-52.2)	15.0 (11.0-20.0)	18.8 (13.2-26.0)	31.7 (20.8-45.0)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Hispanic/Latino  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN102: Percentage of students who strongly agree or agree that their parents or other adults in their family have clear rules and consequences for their behavior	81.7 (72.2-88.4)	76.2 (70.0-81.5)	69.3 (61.1-76.4)	52.7 (31.9-72.6)	Yes
QN103: Percentage of students who went hungry most of the time or always because there was not enough food in their home	4.4 (2.2-8.7)	3.0 (1.6-5.7)	9.6 (3.0-26.5)	20.1 (10.5-35.0)	Yes
QN104: Percentage of students who have ever lived with someone who had a problem with alcohol or drugs	36.8 (20.1-57.5)	26.8 (21.7-32.6)	40.5 (30.7-51.1)	57.9 (36.9-76.4)	Yes
QN105: Percentage of students who have talked with at least one of their parents or guardians about the dangers of tobacco, alcohol, or drug use (during the 12 months before the survey)	62.7 (49.3-74.5)	44.2 (37.5-51.1)	51.5 (41.3-61.7)	36.7 (24.7-50.6)	Yes
QN106: Percentage of students who recall hearing, reading, or seeing a message about avoiding alcohol or other illegal drugs (during the 12 months before the survey)	80.3 (70.8-87.3)	73.8 (67.4-79.4)	71.5 (61.8-79.5)	70.7 (54.5-82.9)	No
QN107: Percentage of students who would most likely talk with their parent or other adult family member about their feelings (among students who report having felt sad, empty, hopeless, angry, or anxious)	25.4 (17.7-35.0)	15.8 (10.9-22.2)	7.6 (3.9-14.0)	11.5 (4.8-25.3)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Hispanic/Latino  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN108: Percentage of students who performed organized community service as a non-paid volunteer (such as serving meals to the elderly, picking up litter, helping out at a hospital, or building homes for the poor, one or more times during the 30 days before the survey)	44.2 (29.8-59.7)	35.1 (28.2-42.7)	24.8 (17.9-33.3)	14.9 (7.8-26.7)	Yes
QN109: Percentage of students who participate in activities run by community groups (including activities run by 4-H, Boys and Girls Club, YMCA, sports clubs, or church groups, for one or more hours during an average week when they are in school)	41.8 (28.7-56.3)	33.8 (26.7-41.8)	20.6 (14.4-28.5)	11.6 (5.9-21.5)	Yes
QN110: Percentage of students who think people are at great risk harming themselves if they smoke one or more packs of cigarettes per day (physically or in other ways)	61.1 (41.6-77.7)	64.6 (58.2-70.6)	49.7 (38.9-60.6)	35.4 (24.3-48.4)	Yes
QN111: Percentage of students who think people are at great risk harming themselves if they have one or two drinks of alcohol nearly every day (physically or in other ways)	27.8 (18.5-39.5)	30.2 (23.4-37.9)	21.4 (14.7-30.1)	16.8 (8.9-29.2)	No
QN112: Percentage of students who think people are at great risk of harming themselves (physically or in other ways) if they have five or more drinks of alcohol once or twice a week (beer, wine, or liquor)	39.4 (26.9-53.4)	39.5 (32.4-47.0)	33.3 (25.5-42.0)	15.6 (8.4-27.2)	No
QN113: Percentage of students who think people are at great risk of harming themselves if they use marijuana once or twice a week (physically or in other ways)	20.3 (13.2-30.0)	19.7 (13.4-28.1)	13.6 (9.0-20.2)	5.7 (2.1-14.7)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Hispanic/Latino  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN114: Percentage of students who think people are at great risk of harming themselves if they take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	56.2 (38.5-72.5)	60.5 (53.4-67.2)	47.0 (37.2-56.9)	33.5 (22.7-46.4)	Yes
QN115: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to smoke tobacco	75.6 (64.7-83.9)	66.6 (59.8-72.7)	49.7 (39.6-59.8)	27.4 (15.9-43.0)	Yes
QN116: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to have one or two drinks of alcoholic beverage nearly every day (beer, wine or liquor)	66.8 (54.1-77.5)	56.1 (49.0-63.1)	36.6 (28.0-46.2)	29.5 (17.4-45.3)	Yes
QN117: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to smoke marijuana	53.6 (38.3-68.2)	37.4 (30.2-45.3)	25.5 (17.9-34.8)	15.7 (8.2-28.0)	Yes
QN118: Percentage of students who reported that their friends feel it would be wrong or very wrong for them to take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	83.3 (74.1-89.7)	74.8 (67.9-80.6)	68.0 (55.8-78.2)	44.6 (25.7-65.1)	Yes
QN119: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to smoke tobacco	94.4 (89.7-97.0)	89.0 (84.6-92.3)	79.6 (65.7-88.8)	61.8 (48.3-73.6)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### New Hampshire High School Survey Risk Behaviors and Academic Achievement Report

**Hispanic/Latino  
Site-Added**

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN120: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to have one or two drinks of an alcoholic beverage nearly every day (beer, wine or liquor)	89.7 (82.3-94.3)	86.1 (81.2-89.9)	78.6 (66.1-87.4)	58.2 (44.7-70.5)	Yes
QN121: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to smoke marijuana	87.8 (80.3-92.7)	78.9 (72.0-84.4)	70.7 (58.3-80.6)	54.4 (41.0-67.1)	Yes
QN122: Percentage of students who reported that their parents feel it would be wrong or very wrong for them to take a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	92.2 (86.1-95.7)	92.6 (88.7-95.2)	88.1 (72.7-95.4)	71.1 (54.3-83.6)	Yes
QN123: Percentage of students who approve or strongly approve of someone their age having one or two drinks of alcohol nearly every day (beer, wine, or liquor)	25.0 (9.2-52.4)	13.6 (9.8-18.7)	14.2 (9.0-21.7)	32.0 (18.1-49.9)	Yes
QN124: Percentage of students who think it would be very easy for them to get some cigarettes if they wanted to	30.2 (19.5-43.5)	39.4 (32.4-46.9)	53.6 (43.6-63.3)	67.8 (50.3-81.4)	Yes
QN125: Percentage of students who think it would be very easy for them to get some alcohol if they wanted to (beer, wine, or liquor)	31.0 (20.9-43.3)	38.8 (32.1-46.0)	43.4 (32.5-55.0)	63.9 (45.9-78.7)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey  
Risk Behaviors and Academic Achievement Report

Hispanic/Latino  
Site-Added

Health Risk Behavior	Percentage of students who engaged in each risk behavior, by type of grades mostly earned in school				Significant Association*
	A's	B's	C's	D's/F's	
	% 95% CI†	% 95% CI	% 95% CI	% 95% CI	
QN126: Percentage of students who think it would be very easy for them to get some marijuana if they wanted to	30.9 (20.7-43.3)	47.2 (40.2-54.3)	61.9 (51.5-71.3)	67.1 (49.7-80.8)	Yes
QN127: Percentage of students who think it would be very easy for them to get some prescription drug without a doctor's prescription if they wanted to (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax)	11.8 (7.2-18.9)	19.4 (14.8-25.0)	22.0 (12.1-36.6)	49.0 (35.9-62.3)	Yes

\*Based on logistic regression analysis controlling for sex, race/ethnicity, and grade in school, p<0.05.

† Confidence interval.